

Product Information

Anti- β -Tubulin antibody, Mouse Monoclonal

Clone AA2, purified from hybridoma cell culture

T8328

Product Description

Anti- β -Tubulin (mouse IgG1 isotype) is derived from the hybridoma AA2 produced by the fusion of mouse myeloma cells and splenocytes from BALB/c mice immunized with purified bovine tubulin. The isotype is determined using a double diffusion immunoassay using Mouse Monoclonal Antibody Isotyping Reagents, Cat. No. ISO2.

Anti- β -Tubulin reacts with human, rat, mouse, and bovine β -tubulin, types I, II, III, and IV. The antibody may be used in various immunochemical techniques including immunoblotting (~ 50 kDa), immunoprecipitation, immunocytochemistry, and immunohistochemistry.¹

Tubulin is the major building block of microtubules. This intracellular, cylindrical, filamentous structure is present in almost all eukaryotic cells. Microtubules function as structural and mobile elements in mitosis, intracellular transport, flagellar movement and the cytoskeleton. Tubulin is a heterodimer that consists of α and β -tubulin. Both subunits have a molecular weight of ~ 50 kDa and share considerable homology. In addition to α -tubulin and β -tubulin, several other tubulins have been identified, bringing the number of distinct tubulin classes to seven. Most of these tubulins have distinct subcellular localization and an emerging diverse set of functions.² Out of the seven different tubulins four new members of the tubulin family were identified, which consist of δ , ζ , η and ϵ -tubulin. η and ϵ -tubulins were discovered by database searches.³ Microtubular systems contain at least three α -tubulin isoforms. Two α -tubulin isoform proteins are very similar but are encoded by two different genes. The third isoform is generated by post-translational modification.⁴ At least three modifications of tubulin subunits have been described: phosphorylation of β -tubulin from brain, removal of the carboxy-terminal tyrosine from α -tubulin in vertebrate tissues, and acetylation of the amino group of lysine(s) in α -tubulin.

Antibodies recognizing α -tubulin, together with monoclonal antibodies to other tubulin types (β -tubulin isotypes I, II, and III, tyrosine tubulin and acetylated- α -tubulin) provide specific and useful tools in studying the intracellular distribution of tubulin, and the static and dynamic aspects of cytoskeleton. In addition, the various tubulin isoforms have been implicated in cancerous processes and are thus suggested as potential targets for cancer therapy.⁵

Reagent

Supplied as a solution in 0.01 M phosphate buffered saline, pH 7.4, containing 15 mM sodium azide as a preservative.

Antibody concentration: ~ 2.0 mg/mL

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For extended storage, freeze at -20 °C in working aliquots. Repeated freezing and thawing, or storage in "frost-free" freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilution samples should be discarded if not used within 12 hours.

Product Profile

Immunoblotting: A working concentration of 1-2 µg/mL is recommended using Neuro-2a cell extract.

Note: In order to obtain the best results using various techniques and preparations, we recommend determining optimal working dilutions by titration.

References

1. Moody, S.A., et al., J. Comp. Neurol., 364, 219-230 (1996).
2. Dutcher, S.K., et al., Curr. Opin. Cell Biol., 13, 49-54 (2001)
3. McKean, P.G., et al., J. Cell Sci., 114, 2723-2733 (2001).
4. LeDizet, M., and Piperno, G., J. Cell Biol., 104, 13-22 (1986).
5. Carlson, R.O., Expert Opin. Investig. Drug, 17, 707-712 (2008).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at [SigmaAldrich.com/techservice](https://www.sigmaaldrich.com/techservice).

Standard Warranty

The applicable warranty for the products listed in this publication may be found at [SigmaAldrich.com/terms](https://www.sigmaaldrich.com/terms).

Contact Information

For the location of the office nearest you, go to [SigmaAldrich.com/offices](https://www.sigmaaldrich.com/offices).

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

MilliporeSigma, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.

© 2022 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

T8328pis Rev 05/22