

Product Information

Anti-Flag® M2 Antibody, Bioreactor grade, Mouse Monoclonal

Clone M2, Purified from Hybridoma Cell Culture

B3111

Product Description

Monoclonal ANTI-FLAG® M2 is a purified immunoglobulin, IgG1, monoclonal antibody, purified from culture supernatant of hybridoma cells, that binds to FLAG® fusion proteins.¹ Unlike ANTI-FLAG® M1 antibody, the M2 antibody will recognize the FLAG® sequence at the N-terminus, Met-N-terminus, C-terminus, or at an internal site of FLAG® fusion proteins. Monoclonal ANTI-FLAG® M2 is useful for identification and capture of FLAG® fusion proteins by common immunological procedures such as Western blots and immunoprecipitation. It is also useful for affinity purification of FLAG® fusion proteins when bound to a solid support.

Monoclonal ANTI-FLAG® M2 binding is not calcium dependent.

Reagent

Supplied as a solution in 0.01 M phosphate buffered saline pH 7.4, containing 15 mM sodium azide as a preservative.

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Safety Data Sheet on the product page online at SigmaAldrich.com for information regarding hazards and safe handling practices.

Storage/Stability

Store the undiluted antibody at -20 °C in working aliquots. Repeated freezing and thawing is not recommended.

Note: Overtime, small amounts of purified antibodies can precipitate from solution due to intermolecular hydrophobic interactions. If a precipitate is observed in this product, briefly centrifuge the vial to pellet the precipitate. Withdraw the desired volume of antibody solution from the clear supernatant for use. This should not alter the performance of the purified antibody in Western blot or immunoprecipitation applications.

Procedure

Improved Western Blot Method for Detecting FLAG® Fusion Proteins using Monoclonal ANTI-FLAG® M2.

1. Separate FLAG® fusion proteins from sample lysates using a standard sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) protocol. Load 2.5–10 µg of total lysate protein per lane.
2. Transfer proteins from the gel to an Immobilon®-P or other polyvinylidene difluoride (PVDF) membrane. Nitrocellulose membranes can be used, but typically result in less sensitivity.
3. Wash the blot in at least 0.5 mL/cm² of Milli-Q® water for 2–3 minutes with mild agitation.
4. Block the blot with at least 0.5 ml/cm² of Phosphate Buffered Saline (PBS), with 5% nonfat milk, for 60 minutes at room temperature with agitation (about 50–60 rpm).
5. Remove the blocking agent and wash once with 0.5 mL/cm² of PBS.
6. Add Monoclonal ANTI-FLAG® M2 to a final concentration of 10 µg/mL to the blot in at least 0.5 mL/cm² of PBS-TWEEN® 20, with 0.5% nonfat milk and incubate at room temperature for 60 minutes.
Note: Using less Monoclonal ANTI-FLAG® M2 antibody may help to reduce background and cross-reactivity. See the "Troubleshooting Guide."
7. Remove the Monoclonal ANTI-FLAG® M2 solution and wash once with at least 0.5 mL/cm² of PBS.
8. Add Anti-Mouse IgG–Peroxidase (Cat. No. A2304) or equivalent, to at least 0.5 mL/cm² of PBS-TWEEN® 20, with 0.5% nonfat milk. Incubate the blots with shaking at room temperature for 60 minutes.

- Wash the blot eight times for a total of 20 minutes in Phosphate Buffered Saline (PBS), plus 0.05% TWEEN® 20.
- Develop the blots using suitable ECL™ substrate for 1 minute.

Immunofluorescence

Monoclonal ANTI-FLAG® M2 may be used in immunofluorescent procedures. A typical concentration for use is 20 µg/mL.²

Product Profile

Protein concentration (E_{280}): 3.0 – 5.0 mg/mL.

Antigenic binding site: N-Asp-Tyr-Lys-Asp-Asp-Asp-Lys-C.

Specificity: Monoclonal ANTI-FLAG® M2 detects a single band of protein on a Western blot from HEK 293 cells over-expressing FLAG® proteins, crude cell lysate.

Sensitivity: Monoclonal ANTI-FLAG® M2 detects 2 ng of FLAG-BAP™ fusion protein on a dot blot using chemiluminescent detection.

Note: In order to obtain best results, it is recommended that each individual user determine working dilution by titration assay.

References

- Brizzard, B.L., et al., Immunoaffinity purification of FLAG® epitope-tagged bacterial alkaline phosphatase using a novel monoclonal antibody and peptide elution. *BioTechniques*, 16, 730-735 (1994)
- Ciaccia, A.V., and Price, E.M., IBI FLAG® Epitope, 1, 4-5 (1992)
- Bjerrum, O.J., and Heegaard. N.H.H., CRC Handbook of Immunoblotting of Proteins, Volume I, Technical Descriptions, CRC Press, (1988) p. 229-236
- Dunbar, B.S. (ed.) Protein Blotting: A Practical Approach, IRL Press, NY, p. 67-70 (1994)
- Fortin, A., et al., A 56- to 54-kilodalton non grata signal in immunoblot analysis using the horseradish peroxidase chemiluminescence system. *Biochem. Cell Biol.*, 72, 239-243 (1994)
- Harlow, E., and Lane, D., Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1988, Product Code A2926
- Layton, Curtis J., Peter L. McMahon, and William J. Greenleaf. "Large-scale, quantitative protein assays on a high-throughput DNA sequencing chip." *Molecular cell* 73.5 (2019): 1075-1082.

- Bi, Xinyan, and Kun-Lin Yang. "Liquid crystals decorated with linear oligopeptide FLAG® for applications in immunobiosensors." *Biosensors and Bioelectronics* 26.1 (2010): 107-111.
- Wang, Jianhao, et al. "Resolving antibody-peptide complexes with different ligand stoichiometries reveals a marked affinity enhancement through multivalency." *Talanta* 115 (2013): 394-400.
- Lakamp, Amanda S., and Michel M. Ouellette. "A ssDNA aptamer that blocks the function of the ANTI-FLAG® M2 antibody." *Journal of nucleic acids* 2011 (2011).
- Roosild, Tarmo P., Samantha Castronovo, and Senyon Choe. "Structure of anti-FLAG® M2 Fab domain and its use in the stabilization of engineered membrane proteins." *Acta Crystallographica Section F: Structural Biology and Crystallization Communications* 62.9 (2006): 835-839.

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

Technical Assistance

Visit the tech service page on our web site at SigmaAldrich.com/techservice.

Standard Warranty

The applicable warranty for the products listed in this publication may be found at SigmaAldrich.com/terms.

MilliporeSigma, Millipore, ANTI-FLAG, FLAG, FLAG-BAP, Milli-Q, Immobilon, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.
© 2020 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

MILLIPORE
SIGMA