

Product Information

Pyocyanin, Ready Made Solution from *Pseudomonas aeruginosa*

R9532

Storage Temperature –20 °C


CAS RN 85-66-5

Synonyms: Sanasin, Sanazin, Pyocyanine

Molecular weight: 210.23

Molecular formula: C₁₃H₁₀N₂O

Purity: ≥ 98% (HPLC)



Product Description

Pyocyanin is a blue-green pigment, which belongs to the Phenazine family. It is an electron acceptor, which stimulates redox cycling in bacteria, liver cells, and human epithelial cell lines.^{1,2} Pyocyanin enhances oxidative metabolism, which increases the formation of intracellular reactive oxygen species (ROS) via reduction of NADPH.^{1,3,4}

Pyocyanin also increases the release of the neutrophil chemoattractant interleukin-8 (IL-8) by airway epithelial cells both *in vitro* and *in vivo*. This involves signal transduction pathways that include oxidants, protein tyrosin kinases, and MAP-kinases. IL-8 secretion by these cells is in synergy with inflammatory cytokines.^{1,4,5} Pyocyanin has been shown to accelerate neutrophil apoptosis *in vitro*. Mice infected with a pyocyanin-deficient strain of *P. aeruginosa* showed elevated levels of neutrophils, and neutrophil chemokines and cytokines, as well as compromised bacterial clearance from the lungs compared with mice infected with a wild type strain. This suggests that pyocyanin production by *P. aeruginosa* suppresses the acute inflammatory response by pathogen-driven acceleration of neutrophil apoptosis and by reducing local inflammation, and that this is advantageous for bacterial survival.⁶

Components

The product is supplied as a 5 mg/mL (24 mM) solution in DMSO.

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses.

Storage/Stability

Store the product sealed at –20 °C. Under these conditions the product is stable for at least 2 years.

References

1. Kanthakumar, K., et al., Mechanisms of action of *Pseudomonas aeruginosa* pyocyanin on human ciliary beat *in vitro*. *Infect. Immun.*, **61**: 2848-2853 (1993).
2. Da Silva, G.A., and de Almeida, E.A., Production of yellow-green fluorescent pigment by *Pseudomonas fluorescens*. *Braz. Arch. Biol. Technol.*, **49**: 411-419 (2006).
3. Price-Whelan, A., et al., Pyocyanin alters redox homeostasis and carbon flux through central metabolic pathways in *Pseudomonas aeruginosa* PA14. *J. Bacteriol.*, **189**: 6372-6381 (2007).

4. O'Malley, Y.Q., et al., *Pseudomonas aeruginosa* pyocyanin directly oxidizes glutathione and decreases its levels in airway epithelial cells. *Am. J. Physiol. Lung Cell. Mol. Physiol.*, **287**: L94-L103 (2004).
5. Denning, G.M., et al., *Pseudomonas* pyocyanin increases interleukin-8 expression by human airway epithelial cells. *Infect. Immun.*, **66**: 5777-5784 (1998).
6. Allen, L., et al., Pyocyanin production by *Pseudomonas aeruginosa* induces neutrophil apoptosis and impairs neutrophil-mediated host defenses *in vivo*. *J. Immunol.*, **174**: 3643-3649 (2005).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Standard Warranty

The applicable warranty for the products listed in this publication may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

MilliporeSigma, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. All other trademarks are the property of their respective owners. Detailed information on trademarks is available via publicly accessible resources.
© 2022 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.