

## Product Information

### CAMK1 $\delta$ Active human, recombinant GST-tagged, expressed in Sf9 cells

Catalog Number **C9619**  
Lot Number 019K1566  
Storage Temperature  $-70^{\circ}\text{C}$

Synonyms: RP11-462F15.1; CKLiK; CaM-K1; CaMKID

#### Product Description

CAMK1 $\delta$  or Ca<sup>2+</sup>/calmodulin-dependent kinase I-like kinase (CKLiK) is activated by Ca<sup>2+</sup> and calmodulin, and is detected in CD34+-derived neutrophils and eosinophils, as well as in mature peripheral blood granulocytes.<sup>1</sup> CAMK1 $\delta$  exhibits Ca<sup>2+</sup>/CaM-dependent activity that is enhanced *in vitro* by phosphorylation of Thr<sup>180</sup> by CaM-K kinase (CaM-KK)alpha, consistent with detection of CAMK1 $\delta$ -activating activity in HeLa cells.<sup>2</sup>

This recombinant product was expressed by baculovirus in Sf9 insect cells using an N-terminal GST-tag. The gene accession number is NM 153498. It is supplied in 50 mM Tris-HCl, pH 7.5, with 150 mM NaCl, 0.25 mM DTT, 0.1 mM EGTA, 0.1 mM EDTA, 0.1 mM PMSF, and 25% glycerol.

Molecular mass: ~68 kDa

Purity:  $\geq 70\%$  (SDS-PAGE, see Figure 1)

Specific Activity: 117–159 nmole/min/mg (see Figure 2)


#### Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.


#### Storage/Stability

The product ships on dry ice and storage at  $-70^{\circ}\text{C}$  is recommended. After opening, aliquot into smaller quantities and store at  $-70^{\circ}\text{C}$ . Avoid repeated handling and multiple freeze/thaw cycles.

**Figure 1.**  
SDS-PAGE Gel of Lot Number 019K1566:  
>90% (densitometry)



**Figure 2.**  
Specific Activity of Lot Number 019K1566:  
138 nmole/min/mg



#### Procedure

##### Preparation Instructions

Kinase Assay Buffer – 25 mM MOPS, pH 7.2, 12.5 mM glycerol 2-phosphate, 25 mM MgCl<sub>2</sub>, 5 mM EGTA, and 2 mM EDTA. Just prior to use, add DTT to a final concentration of 0.25 mM.

Kinase Dilution Buffer – Dilute the Kinase Assay Buffer 5-fold with a 50 ng/μl BSA solution.

Kinase Solution – Dilute the Active CAMK1 $\delta$  (0.1  $\mu\text{g}/\mu\text{l}$ ) with Kinase Dilution Buffer to the desired concentration.

**Note:** The lot-specific specific activity plot may be used as a guideline (see Figure 2). It is recommended that the researcher perform a serial dilution of Active CAMK1 $\delta$  kinase for optimal results.

10 mM ATP Stock Solution – Dissolve 55 mg of ATP in 10 ml of Kinase Assay Buffer. Store in 200  $\mu\text{l}$  aliquots at  $-20\text{ }^{\circ}\text{C}$ .

$\gamma$ - $^{32}\text{P}$ -ATP Assay Cocktail (250  $\mu\text{M}$ ) – Combine 5.75 ml of Kinase Assay Buffer, 150  $\mu\text{l}$  of 10 mM ATP Stock Solution, 100  $\mu\text{l}$  of  $\gamma$ - $^{32}\text{P}$ -ATP (1 mCi/100  $\mu\text{l}$ ). Store in 1 ml aliquots at  $-20\text{ }^{\circ}\text{C}$ .

Substrate Solution – Dissolve the synthetic peptide substrate (KKALRRQETVDAL-amide) in water at a final concentration of 1 mg/ml.

1% phosphoric acid solution – Dilute 10 ml of concentrated phosphoric acid to a final volume of 1 L with water.

#### Kinase Assay

This assay involves the use of the  $^{32}\text{P}$  radioisotope. All institutional guidelines regarding the use of radioisotopes should be followed.

1. Thaw the Active CAMK1 $\delta$ , Kinase Assay Buffer, Substrate Solution, and Kinase Dilution Buffer on ice. The  $\gamma$ - $^{32}\text{P}$ -ATP Assay Cocktail may be thawed at room temperature.
2. In a pre-cooled microcentrifuge tube, add the following solutions to a volume of 20  $\mu\text{l}$ :
  - 10  $\mu\text{l}$  of Kinase Solution
  - 7.5  $\mu\text{l}$  of Substrate Solution
  - 2.5  $\mu\text{l}$  of 5 mM  $\text{CaCl}_2$  solution containing 0.75  $\mu\text{g}$  Calmodulin
3. Set up a blank control as outlined in step 2, substituting 7.5  $\mu\text{l}$  of cold water ( $4\text{ }^{\circ}\text{C}$ ) for the Substrate Solution.
4. Initiate each reaction with the addition of 5  $\mu\text{l}$  of the  $\gamma$ - $^{32}\text{P}$ -ATP Assay Cocktail, bringing the final reaction volume to 25  $\mu\text{l}$ . Incubate the mixture in a water bath at  $30\text{ }^{\circ}\text{C}$  for 15 minutes.
5. After the 15 minute incubation, stop the reaction by spotting 20  $\mu\text{l}$  of the reaction mixture onto an individually pre-cut strip of phosphocellulose P81 paper.

6. Air dry the pre-cut P81 strip and sequentially wash in the 1% phosphoric acid solution with constant gentle stirring. It is recommended the strips be washed a total of 3 times of  $\sim 10$  minutes each.
7. Set up a radioactive control to measure the total  $\gamma$ - $^{32}\text{P}$ -ATP counts introduced into the reaction. Spot 5  $\mu\text{l}$  of the  $\gamma$ - $^{32}\text{P}$ -ATP Assay Cocktail on a pre-cut P81 strip. Dry the sample for 2 minutes and read the counts. Do not wash this sample.
8. Count the radioactivity on the P81 paper in the presence of scintillation fluid in a scintillation counter.
9. Determine the corrected cpm by subtracting the blank control value (see step 3) from each sample and calculate the kinase specific activity

#### Calculations:

1. Specific Radioactivity (SR) of ATP (cpm/nmole)

$$\text{SR} = \frac{\text{cpm of } 5\ \mu\text{l of } \gamma\text{-}^{32}\text{P}\text{-ATP Assay Cocktail}}{\text{nmole of ATP}}$$

cpm – value from control (step 7)  
nmole – 1.25 nmole (5  $\mu\text{l}$  of 250  $\mu\text{M}$  ATP Assay Cocktail)

2. Specific Kinase Activity (SA) (nmole/min/mg)

$$\text{nmole/min/mg} = \frac{\Delta\text{cpm} \times (25/20)}{\text{SR} \times \text{E} \times \text{T}}$$

SR = specific radioactivity of the ATP (cpm/nmole ATP)

$\Delta\text{cpm}$  = cpm of the sample – cpm of the blank (step 3)

25 = total reaction volume

20 = spot volume

T = reaction time (minutes)

E = amount of enzyme (mg)

#### References

1. Verploegen, S. et al., Characterization of the role of CaMKI-like kinase (CKLiK) in human granulocyte function. *Blood*, **106**, 1076-1083 (2005).
2. Ishikawa, Y. et al., Identification and characterization of novel components of a  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase cascade in HeLa cells. *FEBS Lett.*, **550**, 57-63 (2003).

TLD, MAM 02/09-1

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.