Supelco_®

Product Information

BCIP®/NBT Solution, premixed

B6404

Product Description

5-bromo-4-chloro-3-indolyl phosphate (BCIP®) and nitro blue tetrazolium (NBT) are reagents which are widely used in tandem to detect alkaline phosphatase conjugates. This premixed BCIP®/NBT solution is a substrate solution that is designed to visualize alkaline phosphatase conjugates in Western blotting. Several dissertations I have cited use of product B6404 in their protocols.

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Reagent

This premixed BCIP®/NBT solution contains 0.48 mM NBT, 0.56 mM BCIP®, 10 mM Tris HCl (pH \sim 9.2), and 59.3 mM MgCl₂.

Additional Reagents Required (Not provided)

- 10× Tris Buffered Saline (Cat. No. T5912)
- TWEEN® 20 (Cat. No. P2287)
- BSA Fraction V powder (Cat. No. A9647)
- Nonfat-dried milk (Cat. No. M7409)

Storage/Stability

This product may be stored at room temperature.

Procedure

- 1. After the gel is transferred onto a blotting membrane, wash the membrane for 5 minutes with the Washing Solution (see Table 1).
- Incubate the rinsed membrane with primary antibody diluted in Blocking Solution (see Table 1) for 2 hours at room temperature with gentle agitation. Use a dilution based on the manufacturer's recommendation. A blocking step prior to Step 2 is usually not necessary.

- 3. Wash the membrane for 5 minutes with the Washing Solution.
- 4. Incubate the washed membrane with a secondary antibody-alkaline phosphatase conjugate in blocking solution for 2 hours at room temperature with gentle agitation. (A 1:1000 dilution of antibody in blocking solution is recommended.)
- 5. Wash the membrane 3 times for 5 minutes each in Washing Solution.
- Cover the membrane with the premixed BCIP®/NBT solution for 1-5 minutes at room temperature until the desired color is obtained. Use ~20 mL for a 10 × 10 cm membrane.
- 7. The color development can be stopped either:
 (a) by extensive washing with water, or
 (b) by rinsing with a 1% acetic acid solution.

Table 1. Commonly used washing and blocking solutions

Condition	Washing Solution	Blocking Solution
1	1× TBS (20 mM Tris-HCl, pH 7.5, and 0.9% NaCl)	Washing solution + 0.5% TWEEN® 20
2	1× TBS + 0.05% TWEEN® 20	Washing solution + 1-5% BSA
3	1× TBS + 0.05% TWEEN® 20	Washing solution + 2-4% nonfat-dried milk

Selection depends on the stringency required:

- Condition 1 is the least stringent.
- Condition 3 is most stringent.

1

 A version of Condition 3, using Washing Solution 3 and Blocking Solution 3 with 2% nonfat-dried milk, is recommended for most applications.



References

- Blake, M.S. et al., Anal. Biochem., 136(1), 175-179 (1984).
- Horowitz, J.P. et al., J. Med. Chem., 9(3), 447 (1966).
- 3. Smejkal, G.B., and Kaul, C.A., *J. Histochem. Cytochem.*, **49(9)**, 1189-1190 (2001).
- 4. Riemenschneider, Leif, "Enzyme Assisted Nanolithography". University of Bremen, Ph.D. dissertation, p. 67 (2004).
- Hasenbrink-Martini, Guido, "Aufbau eines pharma-/toxikologischen Testsystems für die Säuger-Kaliumkanäle mKir2.1 und hERG mit Hilfe des Modellorganismus Saccharomyces cerevisiae" ("Development of a pharma-/toxicological test system for the mammalian potassium channels mKir2.1 and hERG using the model organism Saccharomyces cerevisiae"). Rheinischen Friedrich-Wilhelms-Universität Bonn, Dr. rer. nat. dissertation, p. 44 (2007).
- Saputhanthri, Pradeepika S., "Characterization of an Arabidopsis thaliana mutant purple patch inflorescence: a novel mutant allele of the COP9 Signalosome Subunit 5". University of Bath, Ph.D. dissertation, p. 48 (2007).
- 7. Jones, Matthew A., "Structure-function analysis of phototropin receptor kinases". University of Glasgow, Ph.D. dissertation, p. 45 (2008).
- 8. Moreno, Pedro Martinez, "Estudio de las propiedades de acetilcolinesterasa y butirilcolinesterasa en el tejido pulmonar humano y sus alteraciones en distintos tipos histológicos de cáncer de pulmón de células no pequeñas" ("Study of the properties of acetylcholinesterase and butyrylcholinesterase in human lung tissue and their alterations in different histological types of non-small cell lung cancer"). Universdad de Murcia, Ph.D. dissertation, p. 69 (2012).
- 9. Makadiya, Nirajkumar, "Functional Characterization of 100K Protein of Bovine Adenovirus Type 3". University of Saskatchewan, Ph.D. dissertation, p. 44 (2013).
- 10. Murphy, Valerie, "Expression, Purification and Characterization of Carbohydrate-Binding Proteins". National University of Ireland, Galway, Ph.D. dissertation, p. 40 (2014).

- Withers, Catherine Nicole Kaminski, "Rad GTPase: Identification of Novel Regulatory Mechanisms and a New Function in Modulation of Bone Density and Marrow Adiposity". University of Kentucky, Ph.D. dissertation, p. 59 (2017).
- 12. Fuchs, Lumila Ivana, "Tricomonosis bovina: caracterización de cepas prevalentes en la provincia de La Pampa e inmunoprofilaxis de la enfermedad mediante el empleo de vacunas experimentales en vaquillonas" ("Bovine trichomoniasis: characterization of prevalent strains in the province of La Pampa and immunoprophylaxis of the disease through the use of experimental vaccines in heifers"). Universdad Nacional de la Plata, Ph.D. dissertation, p. 105 (2017).
- Fernández, Lucia Anaís Torres, "Unraveling TRIM71-mediated RNA regulatory mechanisms during developmental and oncogenic processes". Rheinischen Friedrich-Wilhelms-Universität Bonn, Dr. rer. nat. dissertation, p. 28 (2019).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Terms and Conditions of Sale

Warranty, use restrictions, and other conditions of sale may be found at <u>SigmaAldrich.com/terms</u>.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

