

ProductInformation

L-Histidine monohydrochloride monohydrate, from non-animal source Cell culture tested, meets EP testing specifications

Product Number **H5659**
Store at Room Temperature

Product Description

Molecular Formula: $C_6H_9N_3O_2 \cdot HCl \cdot H_2O$
Molecular Weight: 209.6
CAS Number: 5934-29-2
pI: 7.64¹
pK_a: 1.80 (COOH), 9.33 (NH₂),
6.04 (imidazole group)¹
Specific Rotation: +8.9° to +9.5° (11 mg/ml, 6 M HCl)
Synonyms: (S)- α -amino-1H-imidazole-4-propanoic acid hydrochloride, glyoxaline-5-alanine hydrochloride, His, H

This product is cell culture tested (0.42 mg/ml) and is tested for endotoxin levels.

The essential amino acid L-histidine is one of the three amino acids with a basic side chain, and is very hydrophilic in character. It contains an imidazole group in the side chain. Because the pK_A of the imidazole group is close to physiological pH, the imidazole moiety can be either uncharged or positively charged, depending on the local environment. This property makes histidine residues of exceptional interest in the active sites of many proteins, such as hemoglobin and myoglobin, where the imidazole ring can readily alternate between the charged (imidazolium) and uncharged (imidazole) states to participate in bond formation and breakage. Histidine is biosynthesized from ATP, 5-phosphoribosyl-1-pyrophosphate (PRPP), and glutamine. In turn, histidine is degraded to glutamate by histidase, urocanase, and imidazolonepropionase via the formation of urocanate, 4-imidazolone 5-propionate, and N-formiminoglutamate.^{3,4}

L-Histidine has been used to study cultures of the human T-lymphoblastic leukemia cell line MOLT-4 to study modulation of apoptosis.⁵ Exogenous histidine has been shown to enhance the biosynthesis of lovastatin by cultured *Aspergillus terreus*.⁶ Histidine has been utilized as a single nitrogen source to probe

swarming in *Pseudomonas aeruginosa* on agar.⁷ An *in vivo* study has used L-histidine to diminish the net secretory response of the small intestine of cholera toxin-challenged mice.⁸

Precautions and Disclaimer

For Laboratory Use Only. Not for drug, household or other uses.

Preparation Instructions

This product is soluble in water (50 mg/ml), yielding a clear, colorless solution.

References

1. Molecular Biology LabFax, Brown, T. A., ed., BIOS Scientific Publishers Ltd. (Oxford, UK: 1991), p. 29.
2. The Merck Index, 12th ed., Entry# 9929.
3. Biochemistry, 3rd ed., Stryer, L., W. H. Freeman (New York, NY: 1988), pp. 19-20, 505, 587-588.
4. Textbook of Biochemistry with Clinical Correlations, 5th ed., Devlin, T. M., ed., Wiley-Liss (New York, NY: 2002), pp. 97, 395, 816-817.
5. Franek, F., et al., Antiapoptotic and proapoptotic action of various amino acids and analogs in starving MOLT-4 cells. Arch. Biochem. Biophys., **398(1)**, 141-146 (2002).
6. Hajjaj, H., et al., Lovastatin biosynthesis by *Aspergillus terreus* in a chemically defined medium. Appl. Environ. Microbiol., **67(6)**, 2596-2602 (2001).
7. Kohler, T., et al., Swarming of *Pseudomonas aeruginosa* is dependent on cell-to-cell signaling and requires flagella and pili. J. Bacteriol., **182(21)**, 5990-5996 (2000).
8. Peterson, J. W., et al., Cholera toxin-induced PGE₂ activity is reduced by chemical reaction with L-histidine. Biochim. Biophys. Acta, **1537(1)**, 27-41 (2001).

GCY/RXR 3/07

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.