



## KREBS-RINGER BICARBONATE BUFFER

With 1800 mg/L Glucose, Without Calcium Chloride and Sodium Bicarbonate  
Product Number **K4002**

### Product Description

Although there have been many modifications to the original formulas in efforts to produce fully defined media, salt solutions still play an important role in tissue culture. A salt solution's basic function, to maintain the pH and osmotic balance in the medium and to provide the cells with water and essential inorganic ions, is as valuable today as when it was first developed a century ago

| Components                             | g/L    |
|--|--------|
| D-Glucose                              | 1.8    |
| Magnesium Chloride [Anhydrous]         | 0.0468 |
| Potassium Chloride                     | 0.34   |
| Sodium Chloride                        | 7.0    |
| Sodium Phosphate Dibasic [Anhydrous]   | 0.1    |
| Sodium Phosphate Monobasic [Anhydrous] | 0.18   |

### Precautions and Disclaimer

REAGENT

For R&D use only. Not for drug, household or other uses.

### Preparation Instructions

Powdered salts are hygroscopic and should be protected from moisture. The entire contents of each package should be used immediately after opening. Preparing a concentrated salt solution is not recommended as precipitates may form. Supplements can be added prior to filtration or introduced aseptically to sterile salt solution.

1. Measure out 90% of final required volume of water. Water temperature should be 15-20 °C.
2. While gently stirring the water, add the powdered medium. Stir until dissolved. Do NOT heat.
3. Rinse original package with a small amount of water to remove all traces of powder. Add to solution in step 2.
4. To the solution in step 3, add 1.26 g sodium bicarbonate or 16.8 ml of sodium bicarbonate solution [7.5%w/v] for each liter of final volume of medium being prepared. Stir until dissolved.

5. While stirring, adjust the pH of the medium to 0.1-0.3 pH units below the desired pH since it may rise during filtration. The use of 1N HCl or 1N NaOH is recommended.
6. Add additional water to bring the solution to final volume.
7. Sterilize immediately by filtration using a membrane with a porosity of 0.22 microns.
8. Aseptically dispense medium into sterile container.

### Storage and Stability

Store the dry powdered salts at 2-8 °C under dry conditions and liquid medium at 2-8 °C in the dark. Deterioration of the powdered medium may be recognized by any or all of the following: [1] color change, [2] granulation/clumping, [3] insolubility. Deterioration of the liquid medium may be recognized by any or all of the following: [1] pH change, [2] precipitate or particulates, [3] cloudy appearance [4] color change. The nature of supplements added may affect storage conditions and shelf life of the medium. Product label bears expiration date.

### Procedure

Materials Required but Not Provided

Water for tissue culture [W3500]  
Sodium Bicarbonate [S5761] or  
Sodium Bicarbonate Solution, 7.5% [S8761]  
1N Hydrochloric Acid [H9892]  
1N Sodium Hydroxide [S2770]  
Medium additives as required

### Reference

1. Krebs, H. A. and Henseleit, K. (1932). Untersuchungen über die Harnstoffbildung im Tierkörper in Hoppe-Seyler's Zeitschrift für Physiol. Chemie, 210, 33-66.

Revised: June 2007

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.