

# Novabiochem<sup>®</sup>

## Guide to selection of building blocks













## Fmoc-His(Clt)-OH

The use of Fmoc-His(Trt)-OH for the production of protected peptide fragments can result in the formation of a mixture of fully and partially protected peptides because loss of the  $\tau$ -trityl group can occur during the HFIP/DCM or 1% TFA in DCM treatments used to release products from 2-chlorotrityl or HMPB-AM resins. In contrast, the  $\tau$ -Clt group in Fmoc-His(Clt)-OH is completely stable to these mild acids, enabling the preparation of protected peptides free from partially deprotected by-products.

Furthermore, in model studies we have found Fmoc-His(Clt)-OH to give about a third less enantiomerization than Fmoc-His(Trt)-OH.

## Fmoc-His(Boc)-OH • CHA

Like Fmoc-His(Trt)-OH, the Boc-group protects the  $\tau$ -nitrogen. However, the smaller bulk of the Boc group compared to Trt, coupled with the greater stability of Trt towards nucleophiles, suggests that Boc will be less effective than Trt in suppressing enantiomerization [19]. Several side reactions leading to a C-terminal truncated sequence and the formation of succinimide from Asp-His were encountered when using the above-mentioned derivative on a continuous flow instrument during the synthesis of an N-acetylated undecapeptide [95].

### Related products

853008	Boc-His(Dnp)-OH · isopropanol
853041	Boc-His(Tos)-OH
852052	Fmoc-His(Boc)-OH • CHA
852371	Fmoc-His(Clt)-OH
852211	Fmoc-His(Mtt)-OH
852032	Fmoc-His(Trt)-OH
852161	Fmoc-D-His(Trt)-OH

## 1.6 Methionine

The principal side reaction with methionine is the acid catalyzed oxidation of Met to Met(O) during cleavage and deprotection [96, 97]. In Fmoc chemistry Met is introduced without side-chain protection. The formation of Met(O) is prevented by using ethylmethylsulfide [98] or thioanisole [99] during cleavage and deprotection.

In Boc chemistry, methionine can be introduced in its unprotected form. However, spontaneous oxidation can occur during cleavage with strong acids. Alternatively methionine can be introduced as Met(O) during the synthesis and then reduced to Met using the low-high HF (or TFMSA) cleavage procedure outlined in Method 3-17, page 3.25, Novabiochem Catalog.

Alkylation of methionine during HF cleavage can be prevented by using DMS in combination with *p*-thiocresol or anisole [100]. Met(O) can also be reduced to Met after cleavage using N-methylmercaptoacetamide (MMA)[101-103], ammonium iodide/DMS [104], ammonium fluoride/2-mercaptoethanol [105], TiCl<sub>4</sub> [106] or TMSBr/EDT [58, 107]. Reduction with ammonium iodide/DMS has been shown not to affect disulfide bonds [108].

### Related products

853054	Boc-Met-OH
853093	Boc-D-Met-OH
853035	Boc-Met(O)-OH
852002	Fmoc-Met-OH
852140	Fmoc-D-Met-OH
852225	Fmoc-Met-OPfp
852054	Fmoc-Met(O)-OH
852212	Fmoc-Met(O) <sub>2</sub> -OH

## 1.7 Tryptophan

There are two main side reactions which occur with tryptophan residues: oxidation of Trp during synthesis and alkylation of the indole ring by carbonium ions generated during cleavage [1]. In Boc chemistry, the Trp(For) derivative has been used to overcome both of these problems [1]. The formyl group is stable to acid cleavage reagents and can be deprotected either before normal HF cleavage or by thiolytic deprotection using the low-high HF or TFMSA procedure [109]. These procedures are provided in Method 3-19, page 3.25, and Method 3-22, page 3.27, Novabiochem Catalog. Avoid the use of thioanisole during HF cleavage if your peptide contains tryptophan. In Fmoc chemistry, the Trp has been incorporated without side-chain protection. Problems due to *t*-butylation were normally controlled using scavenging mixtures containing EDT [110-112]. However, sulfonation by the by-products from the deprotection of Mtr [113], Pmc [24] and Pbf [28] protected Arg residues, and reattachment of the peptide to the resin support *via* the side chain of a C-terminal Trp residue [114] can only be reduced, but not eliminated, by varying the scavenger mixtures. These problems were resolved with the introduction by Novabiochem's own research group of Fmoc-Trp(Boc)-OH [25]. This derivative has proven to be extremely effective in preventing these side reactions, and is now firmly established in the repertoire of standard reagents for Fmoc SPPS.

### Mechanism of Boc removal

During normal deprotection with TFA the *t*-butyl moiety is removed, leaving the indole protected with an N-carboxy group [115], thereby preventing alkylation, peptide re-attachment or sulfonation. Tryptophan is subsequently regenerated during the course of the normal lyophilization procedure, since this carbamic acid intermediate readily decarboxylates in acidic aqueous media (Figure 2).




Fig. 2: Mechanism of Boc removal.

### Cleavage of Trp and Arg-containing peptides

N<sup>n</sup>-Boc protection prevents sulfonation of Trp, even with sequences containing multiple Arg residues [116] (Application 1). It is especially effective when used in conjunction with thiol-containing scavenger mixtures such as Reagent K [117], but also works well with silanes [28, 116]. The efficacy of using Trp(Boc) in Fmoc synthesis has now been confirmed by a number of independent studies [28, 118, 119]. The combination of Trp(Boc) and Arg(Pbf) seems to be optimal [28].

## Application 1: Synthesis of peptides containing multiple Arg(Pmc) residues with Trp(Boc)

The peptides shown in Figure 3 were prepared automatically on a NovaSyn® Crystal using either N<sup>trt</sup>-Boc protected or unprotected tryptophan. Coupling reactions were carried out using a mixture of Fmoc amino acid/HOBt/PyBOP® (1:1:1) activated with DIPEA (2 eq.) or Fmoc amino acid pentafluorophenyl esters. Peptidyl resins were cleaved by treatment with TFA/thioanisole/phenol/EDT/water (87.5:5:2.5:2.5:2.5).





Fig. 3: HPLC profiles of total crude products obtained from syntheses of model peptides containing Trp and Arg residues.

## Reduction of Trp by silanes

Trialkylsilanes are very effective scavengers, especially for those highly stabilized cations which are not irreversibly scavenged by thiols, such as trityl [120], Tmob [121] and the Rink linker. However, these reagents can cause reduction of tryptophan residues [122], converting the side-chain indole to an indoline. This effect is greatly reduced with Trp(Boc), as demonstrated in Application 2.

## Peptide re-attachment

This can be a particular problem for peptides containing C-terminal Trp, giving severely reduced yields in some cases. Trp(Boc) is effective in limiting this side reaction.

## Application 2: Cleavage with silanes

The peptide was prepared automatically on a NovaSyn® Crystal using either N<sup>trt</sup>-Boc protected or unprotected tryptophan. Coupling reactions were carried out using a mixture of Fmoc amino acid/HOBt/PyBOP® (1:1:1) activated with DIPEA (2 eq.). Peptidyl resins were cleaved by treatment with TFA/TES/water (90:5:5). The HPLC elution profile of the crude product is shown in Figure 4.




Fig. 4: Cleavage of a model peptide with TFA mixtures containing silanes.

## Related products

853038	Boc-Trp-OH
853086	Boc-D-Trp-OH
853078	Boc-Trp(Boc)-OH
853022	Boc-Trp(For)-OH
853108	Boc-D-Trp(For)-OH
852207	Fmoc-Trp-OH
852150	Fmoc-D-Trp-OH
852050	Fmoc-Trp(Boc)-OH
852164	Fmoc-D-Trp(Boc)-OH
852131	Fmoc-Trp(Boc)-OPfp

## References

- J.M. Stewart & J.D. Young, "Solid phase peptide synthesis", Pierce Chemical Company, Rockford, 198
- G. Barany, et al. (1987) *Int. J. Peptide Protein Res.*, 30, 705.
- G. Fields & R.L. Noble (1990) *Int. J. Peptide Protein Res.*, 35, 161.  
E. Atherton & R.C. Sheppard, "Solid phase peptide synthesis: a practical approach", IRL Press, Oxford, 1989.
- W. C. Chan & P. D. White, "Fmoc solid phase peptide synthesis: a practical approach", Oxford University Press, Oxford, 2000.
- J. Meienhofer in "Chemistry & Biochemistry Amino Acids", G.C. Barrett (Ed.), Chapman & Hall, London, 1985, pp. 297.
- B. Rzesotarska, E. Masukiewicz (1988) *Org. Prep. Proc. Int.*, 427.
- T. Copeland & S. Oroszlan (1988) *Gen. Anal. Techn.*, 5, 109.
- E. Wünsch in "Houben-Weyls Methoden der Organischen Chemie"; E. Müller (Ed.), Vol. 15, Parts 1 & 2, Thieme, Stuttgart, 197
- F.C. Gronvald, et al. in "Peptides 1980", K. Brunfeldt (Ed.), 1981, pp. 111.
- E. Atherton, et al. (1985) *J. Chem. Soc. Perkin Trans. I*, 2065.
- H. Rink, et al. (1984) *Tetrahedron Lett.*, 25, 621.
- I. L. Rodionov, et al. in "Innovations & Perspectives in Solid Phase Synthesis, 3rd International Symposium", R. Epton (Ed.), Mayflower Worldwide Ltd., Birmingham, 1994, pp 669.
- A.S. Verdini, et al. (1992) *Tetrahedron Lett.*, 33, 6541.
- R.H. Mazur (1968) *Experientia*, 24, 661.
- Y. Kiso, et al. (1980) *J. Chem. Soc., Chem. Commun.*, 1063.
- M. Fujino, et al. (1981) *Chem. Pharm. Bull.*, 29, 2825.
- J. Yajima, et al. (1978) *Chem. Pharm. Bull.*, 26, 3752.




19. E. Atherton, et al. in "Peptides 1984", U. Ragnarsson (Ed.), 1985, pp. 153.
20. a) N. Fujii, et al. (1987) *Chem. Pharm. Bull.*, **35**, 3880; b) L. Guo, et al. (1988) *Chem. Pharm. Bull.*, **36**, 4989.
21. A. Penrose & P. D. White, Poster 487 presented at the 19th Meeting of the Federation of European Biochemical Societies, Rome, 1989.
22. R. Ramage & J. Green (1987) *Tetrahedron Lett.*, **28**, 2287.
23. J. Green, et al. (1988) *Tetrahedron Lett.*, **29**, 4341.
24. B. Riniker & A. Hartmann in "Peptides, Chemistry, Structure & Biology, Proc. 11th American Peptide Symposium", J.E. Rivier & G.R. Marshall (Eds.), ESCOM, Leiden, 1990, pp. 950.
25. P. White in "Peptides, Chemistry, Structure & Biology, Proc. 12th American Peptide Symposium", J.A. Smith & J.E. Rivier (Eds.), ESCOM, Leiden, 1992, pp. 537.
26. R. Ramage, et al. (1991) *Tetrahedron*, **47**, 6353.
27. L.A. Carpino, et al. (1993) *Tetrahedron Lett.*, **34**, 7829.
28. C.G. Fields & G.B. Fields (1993) *Tetrahedron Lett.*, **34**, 6661.
29. E. Jaeger, et al. (1993) *Biol. Chem. Hoppe Seyler*, **374**, 349.
30. V. Caciagli & A. S. Verdini in "Peptide Chemistry 1987", T. Shiba & S. Sakakibara (Eds.), Protein Research Foundation, Osaka, 1988, pp. 283.
31. M. Bernatowicz & G. Matsueda in "Peptides, Chemistry, Structure & Biology, Proc. 13th American Peptide Symposium", R.S. Hodges & J.A. Smith (Eds.), ESCOM, Leiden, 1994, pp. 107.
32. D.T. Gish, et al. (1971) *J. Am. Chem. Soc.*, **80**, 2558.
33. H. Gausepohl, et al. in "Peptides 1988, Proc. 20th European Peptide Symposium", G. Jung & E. Bayer (Eds.), Walter de Gruyter, Berlin, 1989, pp. 241.
34. H. Gausepohl, et al. (1989) *Int. J. Peptide Protein Res.*, **34**, 287.
35. P. Rovero, et al. (1993) *Tetrahedron Lett.*, **34**, 2199.
36. G. Barany & R.B. Merrifield in "Peptides 2", E. Gross & J. Meienhofer (Eds.); Academic Press, New York, 1980, pp. 1.
37. S. Mojsov, et al. (1980) *J. Org. Chem.*, **45**, 555.
38. M. Bodanszky & V. du Vigneaud (1959) *J. Am. Chem. Soc.*, **81**, 5688.
39. P. Sieber & B. Riniker in "Innovations & Perspectives in Solid Phase Synthesis, 1st International Symposium", R. Epton (Ed.), SPCC (UK) Ltd., Birmingham, 1990, pp. 577.
40. P. Sieber & B. Riniker (1991) *Tetrahedron Lett.*, **32**, 739.
41. M. Friede, et al. (1992) *Pept. Res.*, **5**, 145.
42. L. A. Carpino, et al. (1995) *J. Org. Chem.*, **60**, 7718.
43. R.D. Dimarchi, et al. (1982) *Int. J. Peptide Protein Res.*, **19**, 88.
44. J.P. Tam, et al. (1988) *Pept. Res.*, **1**, 6.
45. E. Nicolás, et al. (1989) *Tetrahedron Lett.*, **30**, 497.
46. R. Subiros-Funosas, et al. (2011) *Tetrahedron*, **67**, 8595.
47. C.D. Chang, et al. (1980) *Int. J. Peptide Protein Res.*, **15**, 59.
48. Y. Okada, et al. (1987) *J. Chem. Soc., Chem. Commun.*, 1532.
49. R. Dölling, et al. (1994) *J. Chem. Soc., Chem. Commun.*, 853.
50. J. Lauer, et al. (1994) *Let. Pept. Sci.*, **1**, 197.
51. R. Subirós-Funosa, et al. (2011) *Biopolymers*, **98**, 89.
52. Y. Yang, et al. (1994) *Tetrahedron Lett.*, **35**, 9689.
53. a) M. Quibell, et al. (1994) *J. Chem. Soc., Chem. Commun.*, 2343; b) L.C. Packman (1995) *Tetrahedron Lett.*, **36**, 7523.
54. A. Karlström & A. Undén (1996) *Tetrahedron Lett.*, **37**, 4243.
55. J.P. Tam, et al. (1979) *Tetrahedron Lett.*, **42**, 4033.
56. J.M. Stewart, et al. (1979) *J. Med. Chem.*, **45**, 555.
57. J.P. Tam, et al. (1977) *J. Am. Chem. Soc.*, **99**, 7363.
58. F. Albericio, et al. in "Fmoc solid phase peptide synthesis", W. C. Chan & P. D. White (Eds.), Oxford University Press, Oxford, 2000, pp. 77, and references therein.
59. E. Atherton, et al. (1985) *J. Chem. Soc., Perkin Trans. I*, 2065.
60. S.N. McCurdy, Applied Biosystems User Bulletin, Peptide Synthesizer; Applied Biosystems, Foster City, 1989, 25, 42.
61. S.N. McCurdy (1989) *Pept. Res.*, **2**, 147.
62. H. Wang, et al. (2000) *Synth. Commun.*, **30**, 727.
63. A. Chimiak in "Peptides 1962", G.T. Young (Ed.), 1963, pp. 37.
64. H.C. Beyerman in "Peptides 1962", G.T. Young (Ed.), 1963, pp. 53.
65. B. Kamber, et al. (1980) *Helv. Chim. Acta*, **63**, 899.
66. P. Sieber, et al. (1977) *Helv. Chim. Acta*, **60**, 27.
67. H. Lamthanh, et al. (1993) *Int. J. Peptide Protein Res.*, **41**, 85.
68. R. G. Hiskey in "The Peptides, Synthesis, Function & Biology, Vol 3", E. Gross & J. Meienhofer (Eds.), Academic Press, New York, 1981, pp. 157.
69. H. Lamthanh, et al. (1995) *Pept. Res.*, **8**, 316.
70. J. Jezek, et al. in "Peptides 1994, Proc. 23rd European Peptide Symposium", H. Maia (Ed.), ESCOM, Leiden, 1994, pp. 250.
71. K. Akaji, et al. (1993) *J. Am. Chem. Soc.*, **115**, 1138.
72. A. Otaka, et al. (1991) *Tetrahedron Lett.*, **32**, 1223.
73. A. Cuthbertson & B. Indrevoll (2000) *Tetrahedron Lett.*, **41**, 3661.
74. M. Yoshida, et al. (1990) *Chem. Pharm. Bull.*, **38**, 273.
75. H. Tamamura, et al. (1998) *Chem. Commun.*, 151.
76. R. Eritja, et al. (1987) *Tetrahedron Lett.*, **43**, 2675.
77. J. C. Spetzler, et al. (1997) *Let. Pept. Sci.*, **3**, 327.
78. J.P. Tam, et al. (1991) *J. Am. Chem. Soc.*, **113**, 6657.
79. F. Albericio, et al. (1991) *Int. J. Peptide Protein Res.*, **37**, 402.
80. T. M. Postma, et al. (2012) *Org. Lett.*, **14**, 5468; T. M. Postma & F. Albericio (2013) *Org. Lett.*, **15**, 616.
81. K. M. Harris, et al. (2007) *J. Pept. Sci.*, **13**, 81.
82. R. A. D. Bathgate, et al. (2006) *Biochemistry*, **45**, 1043.
83. T. Kaiser, et al. (1996) *Tetrahedron Lett.*, **37**, 1187.
84. Y. X. Han, et al. (1997) *J. Org. Chem.*, **62**, 4307.
85. Y. M. Angell, et al. (2002) *J. Peptide Res.*, **5**, 239.
86. Y. Fujiwara, et al. (1994) *Chem. Pharm. Bull.*, **42**, 72.
87. a) J.H. Jones & W.I. Ramage (1978) *J. Chem. Soc., Chem. Commun.*, 472; b) J.H. Jones & A.V. Stachulski (1979) *J. Chem. Soc., Perkin Trans. 1*, 2261.
88. T. Fuji & S. Sakakibara (1974) *Bull. Chem. Soc. Jpn.*, **47**, 3146.
89. T. Ishiguro & C. Eguchi (1989) *Chem. Pharm. Bull.*, **37**, 506.
90. E. Schnabel, et al. (1968) *Ann. Chem.*, **716**, 175.
91. P. Sieber & B. Riniker (1987) *Tetrahedron Lett.*, **28**, 6031.
92. S. J. Harding, et al. in "Peptides 1994, Proc. 24th European Peptide Symposium", H. Maia (Ed.), ESCOM, Leiden, 1995, pp. 189.
93. M. Mergler, et al. (2001) *J. Pept. Sci.*, **7**, 502.
94. M. Mergler, et al., in "Innovations & Perspectives in Solid Phase Synthesis & Combinatorial Libraries", R. Epton (Ed.), Mayflower Worldwide, Kingswinford, 2002, pp. 235.
95. A. Pessi, et al. (1992) *Int. J. Peptide Protein Res.*, **39**, 58.
96. K. Hofmann, et al. (1965) *J. Am. Chem. Soc.*, **87**, 945.
97. K. Norris, et al. (1971) *Acta. Chem. Scand.*, **25**, 945.
98. S. Guttman & R.A. Boissonnas (1959) *Helv. Chim. Acta*, **42**, 1257.
99. H. Yajima, et al. (1980) *Chem. Pharm. Bull.*, **28**, 121.
100. J. Blake & C.H. Li (1968) *J. Am. Chem. Soc.*, **90**, 5882.
101. R.A. Houghten & C.H. Li in "Peptides, Chemistry, Structure & Biology, Proc. 5th American Peptide Symposium", M. Goodman & J. Meienhofer (Eds.), Pierce Chemical Company, Rockford, 1977, pp. 458.
102. R.A. Houghten & C.H. Li (1978) *Int. J. Peptide Protein Res.*, **11**, 345.
103. R.A. Houghten & C.H. Li (1979) *Anal. Biochem.*, **98**, 36.
104. a) C. Garcia-Echeverria, et al. (1989) *Tetrahedron Lett.*, **30**, 2441; b) H. Huang & D. L. Rabenstein (1999) *J. Pept. Res.*, **53**, 548.
105. L. Guo, et al. (1988) *Chem. Pharm. Bull.*, **36**, 4989.
106. M.W. Pennington & M.E. Byrnes (1995) *Pept. Res.*, **39**.
107. W. Beck & G. Jung (1994) *Let. Pept. Sci.*, **1**, 31.
108. E. Nicolas, et al. in "Innovations & Perspectives in Solid Phase Synthesis, 3rd International Symposium", R. Epton (Ed.), Mayflower Worldwide Ltd., Birmingham, 1994, pp. 619.
109. J.P. Tam, et al. (1983) *J. Am. Chem. Soc.*, **105**, 6443.
110. A. Dryland & R.C. Sheppard (1986) *J. Chem. Soc., Perkin Trans. 1*, 125.
111. C.-D. Chang, et al. (1980) *Int. J. Peptide Protein Res.*, **15**, 485.
112. A.N. Eberle, et al. (1986) *J. Chem. Soc., Perkin Trans. 1*, 361.
113. P. Sieber (1987) *Tetrahedron Lett.*, **28**, 1637.
114. a) E. Atherton, et al. (1988) *Tetrahedron*, **44**, 843; b) J.D. Wade, et al. in "Solid phase Synthesis, 1st International Symposium", R. Epton (Ed.), SPCC (UK) Ltd., Birmingham, 1990, pp. 585.
115. H. Franzen, et al. (1984) *J. Chem. Soc., Chem. Commun.*, 1699.
116. a) P. White, Novabiochem Satellite Meeting, Solid phase Synthesis, 2nd International Symposium, Canterbury, 1991; b) P. White, Novabiochem Innovation Sheet 2/93.
117. D.S. King, et al. (1990) *Int. J. Peptide Protein Res.*, **36**, 255.
118. H. Choi & J.V. Aldrich (1993) *Int. J. Peptide Protein Res.*, **42**, 58.
119. C.G. Fields & G.B. Fields in "Innovations & Perspectives in Solid Phase Synthesis, 3rd International Symposium", R. Epton (Ed.), Mayflower Worldwide Ltd., Birmingham, 1994, pp. 251.
120. D. A. Pearson, et al. (1989) *Tetrahedron Lett.*, **30**, 2739.
121. N.A. Solé & G. Barany (1992) *J. Org. Chem.*, **57**, 5399.
122. B. Kamber & B. Riniker in "Peptides, Chemistry & Biology, Proc. 12th American Peptide Symposium", J.A. Smith & J.E. Rivier (Eds.), ESCOM, Leiden, 1992, pp. 525.


## 2 Selectively protected amino acid building blocks

Novabiochem® has one of the largest collections of orthogonally and quasi-orthogonally protected tri-functional amino acids. These derivatives have many interesting applications in peptide and combinatorial chemistry, such as synthesis of cyclic [1] and branched peptides [2] and the preparation of peptides carrying side-chain modifications [3]; the construction of orthogonal multi-release resins for solution screening of one bead-one compound libraries [4, 5]; the preparation of dendritic or cyclic scaffolds for ligand [6, 7] or protein motif presentation [8, 9]; libraries of cyclic glycopeptides [10]; and synthesis of **b**-turn mimetics [11]. The properties of Novabiochem®'s selectively protected amino acids are given in Table 4, page 11.


### 2.1 Diaminopropionic/butanoic acid, ornithine and lysine derivatives



**Fmoc-Dpr(ivDde)-OH, Fmoc-Dab(ivDde)-OH,  
Fmoc-Orn(ivDde)-OH, Fmoc-Lys(ivDde)-OH, ivDde-Lys(Fmoc)-OH**



**Fmoc-Dpr(Mtt)-OH, Fmoc-Dab(Mtt)-OH,  
Fmoc-Orn(Mtt)-OH, Fmoc-Lys(Mtt)-OH**



#### Fmoc-Lys(Mmt)-OH

The side chains of lysine, ornithine, diaminopropionic acid (Dpr) and diaminobutanoic acid (Dab) are very useful for the introduction of a range of modifications, e.g. biotin, fluorescent labels, etc., as well as for the synthesis of branched and cyclic peptides.

Symmetrically branched peptides or MAP core molecules can be produced using the symmetrically protected Fmoc-Lys(Fmoc)-OH, in which both the  $\alpha$ - and  $\epsilon$ -amino groups are simultaneously deprotected with 20% piperidine [12].

However, for specific modification of side-chain amino groups, the use of selective, or ideally orthogonal, protecting group strategies are required. For such applications Novabiochem is able to offer the following derivatives.

#### Dde/ivDde

Since the introduction of the Dde [2] and ivDde [13] amino-protecting groups in 1993 and 1998, respectively, the Fmoc/xDde strategy has become the standard approach for the synthesis of branched, cyclic and side-chain modified peptides by Fmoc SPPS, with over 200 publications citing the use of these protecting groups [14].

Their utility stems from the fact that Dde and ivDde-protected primary amines are stable to 20% piperidine and TFA but are cleaved with 2% hydrazine in DMF. Thus, amino groups protected by these groups can be selectively unmasked on the solid phase without affecting the side-chain protecting groups of other residues, facilitating subsequent site-specific modification. Furthermore, the reaction can be monitored by spectrophotometry since the indazole cleavage product absorbs strongly at 290 nm (Figure 1).




Fig. 5: Removal of ivDde.

Examples of the use of the Fmoc/Dde strategy include: branched [2] and di-epitopic peptides [15]; MAP core molecules and lipo-MAPs [16 -18]; cyclic peptides [1, 19]; TASP molecules [8]; templates for combinatorial chemistry [6, 7]; synthetic proteins [9]; fluorescently-labeled peptides [3]; modified Lys and Dpr derivatives [20, 21]. the synthesis of multi-functional probes [22]; and ubiquitinated peptides [23].

The choice between Dde or ivDde depends on the application. Dde is easier to remove than ivDde but is much less robust. Dde has been observed to undergo migration during piperidine mediated deprotection of N- $\epsilon$ -Fmoc-Lys and N-terminal Dpr residues, leading to scrambling of its position within the peptide chain; and partial loss has been noted during the synthesis of long sequences [24]. The more hindered ivDde on the otherhand does not undergo leaching or side-chain migration to any significant extent, except in the special case of Dpr [25] (see Method 3), but can occasionally prove extremely hard to remove, particularly if located at the C-terminus of a peptide or in an aggregated region of the sequence.

Fmoc derivatives of lysine are available in which the  $\alpha$ - and  $\epsilon$ -amino groups are protected with Dde and ivDde: Fmoc-Lys(Dde)-OH, Dde-Lys(Fmoc)-OH, Fmoc-Lys(ivDde)-OH; and ivDde-Lys(Fmoc)-OH. For Dpr and Dab, only side-chain ivDde-protected derivatives are available: Fmoc-Dpr(ivDde)-OH and Fmoc-Dab(ivDde)-OH.

As hydrazine can also remove Fmoc groups as well as Dde and ivDde groups, when preparing cyclic or side-chain modified peptides by the Fmoc/Dde strategy, assembly of the peptide backbone is generally completed prior to deprotection of the Dde/ivDde groups. The *N*-terminus of the peptide should be protected with Boc, which can be achieved either by direct incorporation of the *N*-terminal residue as a Boc protected amino acid or acylation of the free *N*-terminal amino group with Boc<sub>2</sub>O.

Removal of Dde or ivDde is achieved by treating the resin with 2% hydrazine in DMF, although in cases where ivDde removal has proved difficult, solutions of as much as 10% hydrazine have been employed. The process can be followed spectrophotometrically at the same wavelength used for monitoring removal of Fmoc, since the reaction-product of the Dde or ivDde group with hydrazine is a chromophoric indazole derivative (Figure 5). Typical traces obtained for Dde removal on the NovaSyn® Crystal and Millipore 9050 peptide synthesizers are shown in Figure 6; ivDde gives similar results. Dde and ivDde also stable to the normal reagents employed for Boc cleavage (TFA or 50% TFA in DCM) and to DBU at the normal concentrations (approx. 2%) used for Fmoc removal.

As previously mentioned, if the ivDde group is close to the C-terminus of the peptide or the peptide has aggregated, removal ivDde can be very sluggish and often incomplete. The problem can be avoided for peptides containing side-chain modified lysine by using ivDde-Lys(Fmoc)-OH instead of Fmoc-Lys(ivDde)-OH for incorporation of those residues. The use of the former allows side-chain modification of lysine to be accomplished during chain extension. Following incorporation of ivDde-Lys(Fmoc)-OH into the peptide chain, the side-chain Fmoc group can be removed with piperidine, the side-chain amino group reacted with the desired carboxylic acid-functionalized moiety, before removal of the ivDde with hydrazine, and chain extension in the usual manner.

Recently, complete orthogonality of Dde with Fmoc has been demonstrated if hydroxylamine hydrochloride/imidazole (1.3:1) in NMP instead of hydrazine in DMF is used for Dde removal [26].

### Method 3: Special procedures for chain extension following introduction of Fmoc-Dpr(ivDde)-OH

#### Fmoc removal

1. Treat the Fmoc-Dpr(ivDde)-peptidyl resin with morpholine-DMF (1:1) (2 x 15 min).
2. Quickly wash resin with DMF.

#### Addition of next amino acid

1. Treat resin with Fmoc-AA-OH (1.5 eq.) and DIPEA (3 eq.) in DMF at -20°C.
2. Add PyBOP (1.6 eq.) to the slurry and allow to warm to rt.

In the worst case (Fmoc-Val), migration observed when using this procedure is <10%.




Fig. 6: Spectroscopic monitoring of the removal of Dde based protecting groups.

### Method 2: Selective removal of Dde/ivDde with 2% hydrazine in DMF

#### Batch

1. Place the peptidyl-resin in a flask and treat with 2% hydrazine monohydrate in DMF (25 ml/g). Stopper the flask and leave to stand at rt for 3 min.
2. Filter the resin and repeat the hydrazine treatment two more times. Wash the partially protected resin with DMF.

#### Continuous flow

1. Flow 2% hydrazine monohydrate in DMF at 3 ml/min through the peptidyl resin packed in a 1 cm diameter reaction column. Deprotection can be followed by monitoring spectrophotometrically at 290 nm the absorbance of the column eluant using a 0.1 mm path length cell.
2. When the reaction is complete, as indicated by return of the absorbance to its original value, flush the column with DMF.

### Mtt

The Mtt group can be removed from the side chain of lysine or ornithine using 1% TFA in DCM (Method 4) [27, 28] or with DCM/HFIP/TFE/TES (6.5:2:1:0.5) (Method 5) [29], allowing selective removal in the presence of other side-chain protecting groups which require up to 95% TFA for removal. When using TFA, addition of 1-5% TIS or MeOH is recommended to quench the trityl cations released. Preliminary results indicate MeOH scavenging prevents loss of t-butyl groups and premature cleavage from Rink Amide resins [30] (Method 4b).

The application of Fmoc-Lys(Mtt)-OH is illustrated by the synthesis of a branched peptide in which the sequence Ile→Leu was assembled on an amide resin (Application 3). The *N*-terminal Ile residue was introduced as the Boc-amino acid. The Mtt group was removed from the ε-amino group of the lysine residue with 1% TFA (Method 4a). The Gly→Glu branch was then assembled on the lysine side chain using normal Fmoc chemistry. The peptide was cleaved with 95% TFA, removing all other side-chain protecting groups, as well as the *N*-α-Boc group from the Ile.

Applications of Mtt derivatives include the use of Lys(Mtt) in conjunction with Lys(Dde) to prepare fluorescence-quenched substrates [3], and of Lys(Mtt) and Orn(Mtt) in the preparation of peptidomimetic macrocycles [11, 31].

Table 4: Protecting groups.

Name	Structure	Removed by	Stable to
O-Ally/Alloc		3 eq. Pd(Ph <sub>3</sub> P) <sub>4</sub> in CHCl <sub>3</sub> /AcOH/NMM (37:2:1)	TFA, piperidine, hydrazine
Azido		3 eq. Me <sub>3</sub> P in dioxane/water	TFA, piperidine
O-tBu		95% TFA	1% TFA, piperidine, Pd(O), hydrazine
STmp		b-Mercaptoethanol, 0.1 NMM in DMF	TFA(partial), piperidine
N-xDde		2% hydrazine in DMF or 20% hydroxylamine/ 15% imidazole in NMP/DCM (5:1)	TFA, piperidine, Pd(0), DBU
O-Dmab		2% hydrazine in DMF	TFA, piperidine, Pd(0), DBU
N-Fmoc		20% piperidine in DMF	TFA, Pd(0)
S-Mmt		1% TFA in DCM containing 1-5% TIS	Pd(0), hydrazine, piperidine
N-Mmt		AcOH/TFE/DCM (1:2:7)	
N-Mtt		1% TFA in DCM containing 1-5% TIS	Pd(0), hydrazine, piperidine
O-2-(PhiPr)		2% TFA in DCM containing 1-5% TIS	Pd(0), piperidine
O-Trt		1% TFA in DCM containing 1-5% TIS	Pd(0), hydrazine, piperidine
O-2-ClTrt		1% TFA in DCM containing 1-5% TIS	Pd(0), hydrazine, piperidine

# Orthogonal protecting groups




Fig. 7: HPLC of crude branched peptide.




Fig. 8: PD-MS of crude product; peaks at 1427.5 and 1444.5 are due to  $M+Na^+$  and  $M+K^+$ , respectively.

### Application 3: Synthesis of H-Ile-Pro-Glu-Lys(H-Gly-Asp-Phe-Glu-Glu)-Glu-Thr-Leu-NH<sub>2</sub>

Boc-Ile-Pro-Glu(OtBu)-Lys(Mtt)-Glu(OtBu)-Thr(tBu)-Leu-NovaSyn® KR was prepared using PyBOP®/HOBt chemistry on a NovaSyn® Crystal fully automatic, continuous flow synthesizer. Lys(Mtt) was deprotected by flowing 1% TFA in DCM through the reaction column and monitoring the removal of Mtt spectrophotometrically at 470 nm.

After neutralization of the TFA salt with piperidine, the synthesis continued in the usual way on the side chain of lysine.

The peptide was cleaved using TFA/water/TIS (95:2.5:2.5) for 3 h. This treatment also removed all side-chain protecting groups as well as the N- $\alpha$ -Boc group on Ile.

The crude product was analyzed by HPLC (Figure 4-7) and PD-MS, expected 1406.5, found 1405.7 (Figure 8).

### Method 4-4: Removal of trityl groups on solid phase

#### Batch-wise method

1. Pre-swell the dry resin (1 g) with DCM in a sintered glass funnel (of a type with a tap and stopper). Remove excess DCM.
2. Add 94:1:5 DCM/TFA/TIS (10 ml), seal funnel and shake for 2 min. Remove solvent by applying N<sub>2</sub> pressure.
3. Repeat step 2 three times.
4. Wash resin with DCM and dry under vacuum.

#### Flow method

1. Pre-swell resin (1 g) with DCM and pack into reaction column.
2. Pump 1% TFA in DCM (2 ml/min) through resin. The reaction can be followed by measuring the absorbance of the column eluant using a 0.1 mm flow cell at 460 nm<sup>3</sup>.
3. Once reaction is finished, as indicated by the absorbance returning to baseline, flush column with DCM.

<sup>3</sup>If the peptide contains other trityl-based protecting groups, the level will not return to baseline owing to slow leaching of Trt groups [27].

### Method 5: Removal of Mtt from Lys on solid phase

#### a) DCM/HFIP/TFE/TES [29]

1. Add DCM/HFIP/TFE/TES (6.5:2:1:0.5) (20 ml/g of resin) to peptidyl resin.
2. Leave mixture to stand with gentle agitation for 1 h.
3. Remove a small sample of resin and wash with DCM. Add 1% TFA in DCM to resin sample. If an orange color is formed instantly then leave the reaction for a further hour.
4. Once the trityl test is negative, wash the resin with DMF, 10% DIPEA in DMF, DMF and use in next step of synthesis.

#### b) TFA/DCM/MeOH [28]

1. Add TFA/DCM/MeOH (1:98:1) (15 mg/ml) to peptidyl resin. Drain after 1 min.
2. Add fresh TFA/DCM/MeOH (1:98:1) (15 mg/ml) to peptidyl resin and leave for 16 h.
3. Wash the resin with DMF, 10% DIPEA in DMF, DMF and use in next step of synthesis.

### Mmt

The Mmt group is considerably easier to remove than Mtt. It is cleaved rapidly from the side-chain of lysine using the same methods that were described for Mtt but can be even removed using AcOH/TFE/DCM (1:2:7) [32] or HOBt in DCM/TFE (Method 5-4). The use of Fmoc-Lys(Mmt)-OH is ideal for those cases where removal of Mtt is problematic. Owing to the extreme acid sensitivity of the Mmt group, coupling of Fmoc-Lys(Mmt)-OH is best carried out using PyBOP®/DIPEA or other base-mediated coupling methods.

### Alloc

The Alloc group is stable to treatment with piperidine and TFA, but can be easily removed under mild conditions by Pd(0) catalyzed allyl transfer [33]. A variety of methods have been used for this step, but perhaps the most useful is that of Kates, *et al.* utilizing Pd(Ph<sub>3</sub>P)<sub>4</sub>/CHCl<sub>3</sub>/HOAc/NMM [34] (Method 7). The Alloc group is not compatible with the conditions employed for removal of ivDde [35]. It is thought that the presence of a small amount of diazine in hydrazine causes reduction of the double bond in the allyl group. Fortunately, this side reaction can be easily overcome by the addition of allyl alcohol to the hydrazine reagent.

### Azido

Azides can be reduced under mild conditions with thiols or phosphines to the corresponding amines. Therefore, Fmoc-protected amino acids bearing side-chain azido functionalities are useful tools for preparing branched and side-chain modified peptides. The azido group is stable to coupling conditions and to treatment with piperidine. It is also stable to TFA cleavage conditions provided thiols are omitted from the cocktail [36]. Reduction of the azide on the solid phase selectively unmasks the side-chain amino group without affecting any other amino-acid residues [37, 38].


### Method 6: Reduction of azido group on solid phase

1. Wash resin with three times with dioxane and dioxane/water (4:1).
2. Drain resin and add 1 M Me<sub>3</sub>P in toluene (6 eq.) in dioxane/water (4:1). Gently agitate for 30 mins.
3. Remove a small sample of resin and wash with dioxane then DCM. Add 95% TFA to leave to cleave for 1.5 h. Analyze by HPLC to check completeness of reduction of azido group.


## Related products

852321	Fmoc- $\gamma$ -azidohomalanine
852326	Fmoc- $\epsilon$ -azidonorleucine
852322	Fmoc- $\delta$ -azidonorvaline
852352	(2S,3S)-Fmoc-Abu(3-N <sub>3</sub> )-OH
852084	Fmoc-Dab(ivDde)-OH
852167	Fmoc-D-Dab(ivDde)-OH
852092	Fmoc-Dab(Mtt)-OH
852083	Fmoc-Dpr(ivDde)-OH
852243	Fmoc-D-Dpr(ivDde)-OH
852089	Fmoc-Dpr(Mtt)-OH
852124	Fmoc-Lys(Alloc)-OH
852057	Fmoc-Lys(Dde)-OH
852147	Fmoc-D-Lys(dde)-OH
854000	Dde-Lys(Fmoc)-OH
852082	Fmoc-Lys(ivDde)-OH
852369	Fmoc-D-Lys(ivDde)-OH
852370	ivDde-Lys(Fmoc)-OH
852065	Fmoc-Lys(Mtt)-OH
852094	Fmoc-Lys(Mmt)-OH
852088	Fmoc-Orn(ivDde)-OH
852075	Fmoc-Orn(Mtt)-OH
852351	cis-Fmoc-Pro(4-N <sub>3</sub> )-OH


## 2.2 Hse, Ser, Thr, Tyr and Cys derivatives




Fmoc-Cys(Mmt)-OH




Fmoc-Hse(Trt)-OH



Fmoc-Ser(Trt)-OH



Fmoc-Thr(Trt)-OH



Fmoc-Tyr(2-ClTrt)-OH

### 2-ClTrt/Trt/Mmt

2-ClTrt, Trt and Mmt groups can be removed from the side chains of Tyr, Hse/Ser/Thr [39] and Cys [40] using only 1% TFA in DCM, leaving all other protecting groups intact (Method 4). This allows the selective

deprotection of a single residue for subsequent side-chain modification. In products with multiple Ser, Thr or Tyr residues, selectivity is possible by incorporating all other such residues, with the exception of the residue to be modified, as the *t*-butyl ether. Trityl group removal is an equilibrium process, so must be performed using silanes to scavenge the trityl cations, or in a continuous flow mode to drive the equilibrium. For Ser(Trt), Coba, *et al.* [41] have described the use of 20% dichloroacetic acid in DCM for 10 min for Trt removal. This procedure may also have wider application for the deprotection of other trityl-protected amino acids.

The most common application of the Ser/Thr/Tyr derivatives is in the preparation of phosphopeptides by the post-synthetic method using a phosphoramidite reagent (section 3.11.3, page 3.41, Novabiochem Catalog).

In a comparative study, purer products were obtained using side-chain trityl protected amino acids than with standard *t*-butyl protected amino acids [42].

### tButhio/STmp

The tButhio/STmp can be removed from the side chain by reduction with thiols [43, 44] (Method 3-41, page 3.34, Novabiochem Catalog).


## Related products

852022	Fmoc-Cys(tButhio)-OH
852031	Fmoc-Cys(Mmt)-OH
852373	Fmoc-Cys(STmp)-OH
852059	Fmoc-Hse(Trt)-OH
852046	Fmoc-Ser(Trt)-OH
852166	Fmoc-D-Ser(Trt)-OH
852066	Fmoc-Thr(Trt)-OH
852241	Fmoc-D-Thr(Trt)-OH
852080	Fmoc-Tyr(2-ClTrt)-OH

## 2.3 Asp and Glu derivatives

Orthogonally and quasi-orthogonally protected Asp and Glu derivatives have numerous applications in peptide synthesis and combinatorial chemistry. *a*-Esters are particularly useful for the preparation of head-to-tail cyclic peptides by on-resin cyclization, whereas selectively protected *b*- and *g*-esters of Asp and Glu can be utilized in the synthesis of glycopeptides and side chain-to-side chain or head-to-side chain lactam bridged peptides [33, 34, 45-48]. Similar strategies can also be applied to the construction of cyclic library templates [6, 7].

### Allyl



### Fmoc-Asp/Glu-OAlI/Fmoc-Asp/Glu(OAlI)-OH

Allyl esters are stable to treatment with piperidine and TFA but can be easily removed under mild conditions by Pd(0) catalyzed allyl transfer [33], as previously described for the Alloc group (Method 7).

The allyl group is not compatible with the conditions employed for removal of ivDde [35]. It is thought that the presence of a small amount of diazine in hydrazine causes reduction of the double bond in the

allyl group. Fortunately, this side reaction can be easily overcome by the addition of allyl alcohol to the hydrazine reagent.

## Method 7: Removal of allyl protecting groups


**IMPORTANT:** This reaction is air-sensitive and all manipulations should be carried out under Ar.

1. Weigh the peptidyl resin into a test tube and dry at 40°C under high vacuum. Seal the tube with a rubber septum. Flush the vessel with a stream of Ar delivered via a needle inserted through the septum.
2. Weigh Pd(PPh<sub>3</sub>)<sub>4</sub> (3 eq.) into a dry test tube, add CHCl<sub>3</sub>-AcOH-N-methylmorpholine (37:2:1) (15 ml/g of resin), dissolve catalyst by bubbling a stream of Ar through the solution, and seal the tube with a rubber septum.
3. Transfer this mixture using an Ar flushed gas-tight syringe to the tube containing the resin. Leave to stand for 2 h with occasional gentle agitation.
4. Transfer the resin to a sintered glass funnel and wash consecutively with 0.5% DIPEA in DMF and sodium diethyldithiocarbamate (0.5% w/w) in DMF to remove the catalyst.


The above procedure can also be carried out on any automated peptide synthesizer which uses N<sub>2</sub> agitation for dissolution and transfer of reagents. The Pd(PPh<sub>3</sub>)<sub>4</sub> catalyst should be weighed out into an amino acid cartridge and dissolved in CHCl<sub>3</sub>-AcOH-N-methylmorpholine (37:2:1) with Ar agitation. The cartridge should be sealed and placed in the instrument autoloader as normal. The instrument should then be programmed to transfer the contents of the vial to reaction vessel or column without the addition of further reagents. If the instrument has any spare solvent reservoirs, these can be filled with 0.5% DIPEA in DMF and sodium diethyldithiocarbamate (0.5% w/w) in DMF to allow automated washing of the resin following allyl deprotection.

**NOTE:** If the N-terminal Fmoc group is removed after cleavage of the allyl ester or if a carbodiimide is to be used to effect cyclization, then the resin should also be washed with HOBt/DMF.

## Dmab



## Fmoc-Asp-ODmab, Fmoc-Glu-ODmab



## Fmoc-Asp(ODmab)-OH, Fmoc-Glu(ODmab)-OH

Dmab esters were developed in collaboration between Prof. B. W. Bycroft and Dr. W. Chan, of the University of Nottingham, and Novabiochem®, to provide a complement to Dde amino protection. The design of the Dmab blocking group is based on the safety-catch principle and the known propensity of *p*-aminobenzyl esters to undergo 1,6-elimination [49, 50]. The safety catch is provided by the ivDde group, which protects the amino function of the unstable *p*-aminobenzyl ester during synthesis.

## Deprotection

Dmab protection is quasi-orthogonal to the Fmoc/tBu strategy, since Dmab esters are stable to 20% piperidine in DMF and to TFA, but are quantitatively cleaved with 2% hydrazine in DMF within minutes. Removal of Dmab involves a two-step process: treatment with hydrazine initially removes the N-ivDde group; this is then followed by collapse of the resultant *p*-amino benzyl ester, with concurrent release of the carboxylic acid (Figure 9).

The deprotection reaction can either be carried out in a batch-wise or continuous flow manner. In the latter case, the reaction can be monitored spectrophotometrically at 290 nm by following release of the indazole by-product. Sluggish cleavage of the aminobenzyl moiety has been occasionally observed [51-54], and appears to be very sequence dependent. In these instances washing the support with 20% DIPEA in DMF/water (90:10) [51] or 5 mM sodium hydroxide in methanol [54] has been found to be efficacious. As hydrazine will remove Fmoc, assembly of the peptide backbone must be completed prior to deprotection of the Dmab side chain. The N-terminus of the peptide should be protected with Boc. This can be achieved either by direct incorporation of the N-terminal residue as a Boc protected amino acid or acylation of the free N-terminal amino group with Boc<sub>2</sub>O.

Fmoc-Asp-ODmab has been employed to prepare a cyclic analog of pyrrolicorin [55], a 29-mer head-to-tail cyclic peptide [56], and chlorofusin peptide [57].




Fig. 9: Removal of Dmab.

## 2-PhiPr



## Fmoc-Glu(O-2-PhiPr)-OH Fmoc-Asp(O-2-PhiPr)-OH

The 2-phenylisopropyl group (2-PhiPr) [58] can be removed from the side chain of Asp and Glu using 1% TFA as in the protocol given in Method 7. The use of this protecting group in combination with Mtt on Lys or Orn provides an excellent strategy for the synthesis of side chain-to-side chain lactam bridged peptides. In contrast to Dmab and All, the 2-PhiPr group offers significant protection against aspartimide formation, which makes Fmoc-Asp(O-2-PhiPr) the derivative of choice for the synthesis of cyclic peptides that are prone to this side reaction.

## Cyclic peptide synthesis

In recent years the synthesis and biological properties of cyclic peptides has attracted considerable interest. Introduction of conformational restraint through head-to-tail cyclization has become a standard strategy in medicinal chemistry for increasing the receptor affinity and selectivity of peptide ligands. Furthermore, cyclization has often been employed as a means of prolonging the duration of action of peptide hormones, since

in general cyclic peptides are more stable to proteolysis than their linear counterparts [59]. Cyclic peptides are also used as synthetic immunogens [60], as by restricting conformational flexibility, the peptide is thought to adopt a conformation which more closely mimics that of the epitope as presented on the surface of the native protein.

There are two approaches currently available for the preparation of head-to-tail cyclic peptides. The first involves cyclization of a protected fragment in solution [61], and the second, on-resin cyclization of a peptide attached *via* the side chain of an Asp or Glu residue [45-48, 49, 62]. The solid phase technique is the more straightforward, as excess reagents and by-products can be easily removed by washing of the resin. In addition, this method often gives superior results due to resin induced pseudo-dilution effects. The synthesis of cyclic peptides has been recently reviewed [63].

## Peptide assembly




Fig. 10: Head-to-tail cyclization using  $\alpha$ -allyl esters.

Figure 10 shows the synthesis of cyclic peptide using Fmoc amino acid allyl esters.

For the production of head-to-tail cyclic peptides, the C-terminal amino acid, either Fmoc-Asp-OX or Fmoc-Glu-OX, is attached to the resin via its side chain. If an Asn or Gln residue is required then an amide resin, such as NovaSyn® TGR, should be employed. Chain elongation can be carried out using standard Fmoc SPPS procedures.

Problems of aspartimide or piperidide formation can occur with both Dmab and allyl [666] esters of aspartic acid, involving the nitrogen of the preceding residue or linker. In the former case, this side reaction can be overcome by employing a N-Hmb protected derivative for introduction of the preceding residue [64, 66].

The yield and purity of peptides obtained by this approach are heavily influenced by the choice of solid support, with higher loading resins giving the poorest results [33]. PEG-PS composite resins, such as NovaSyn® TGA and TGR, are particularly recommended as these supports swell well in the chlorinated solvents used for allyl deprotection and the tentacle nature of the resin matrix helps minimize inter-chain oligomerization.

## Cyclization

Cyclization occurs upon activation of the free resin-bound carboxy group with coupling reagents such as TBTU, PyBOP® or DIPCIDI/Oxyma Pure. The reaction normally takes between 2-5 hours, and can be monitored using TNBS or ninhydrin.




Fig. 11: Synthesis of cyclo(Tyr-D-Trp-Val-D-Trp-D-Trp-Arg-Glu) using a Dmab ester.

NOTE: An excess of uronium coupling reagents (i.e. TBTU, HBTU) should be avoided as this can result in capping of resin amino groups. Furthermore, in common with other fragment condensation reactions, epimerization may accompany activation of the carboxy-terminal residue. The extent of this side reaction will vary depending on the solvent, coupling reagent used and peptide sequence. Some experimentation may be required to optimize reaction conditions. The use of Oxyma Pure/ DIPCIDI activation, for this step is particularly recommended.

The strategy for using Dmab is exemplified through synthesis of a cyclic peptide (Application 4).

### Related products

852072	Fmoc-Asp-OAll
852122	Fmoc-Asp(OAll)-OH
856023	Fmoc-Asp(Wang resin)-OAll
856121	Fmoc-Asp(Wang resin LL)-OAll
852079	Fmoc-Asp-ODmab
852078	Fmoc-Asp(ODmab)-OH
856123	Fmoc-Asp(Wang resin LL)-ODmab
852086	Fmoc-Asp(O-2-PhiPr)-OH
852335	Fmoc-Asp-O-2-PhiPr
852073	Fmoc-Glu-OAll
852123	Fmoc-Glu(OAll)-OH
856024	Fmoc-Glu(Wang resin)-OAll
856022	Fmoc-Glu(Wang resin)-ODmab
852077	Fmoc-Glu-ODmab
852076	Fmoc-Glu(ODmab)-OH
852117	Fmoc-Glu-O-2-PhiPr
852085	Fmoc-Glu(O-2-PhiPr)-OH

### References

- G.B. Bloomberg, et al. (1993) *Tetrahedron Lett.*, **34**, 4709.
- B. W. Bycroft, et al. (1993) *J. Chem. Soc., Chem. Commun.*, 778.
- P. Hoogerhout, et al. (1999) *J. Pept. Res.*, **54**, 436.  
M. Lebl, et al. (1993) *Int. J. Peptide Protein Res.*, **41**, 201.
- J. M. Kerr, et al. (1993) *J. Am. Chem. Soc.*, **115**, 2529.
- J. Eichler, et al. (1994) *Pept. Res.*, **7**, 300.
- J. Eichler, et al. in "Innovation & Perspectives in Solid Phase Synthesis, 4th International Symposium", R. Epton (Ed.), Mayflower Scientific Ltd., Birmingham, 1996, pp. 201.
- P. Dumy, et al. in "Peptides 1994, Proc. 23rd European Peptide Symposium", H. Maia (Ed.), ESCOM, Leiden, 1995, pp. 307.
- C. G. Fields, et al. (1993) *Biopolymers*, **33**, 1695.
- V. Wittmann & S. Seeberger (2000) *Angew. Chem. Int. Ed.*, **39**, 4348.
- a) C. Park & K. Burgess (2001) *J. Comb. Chem.*, **3**, 257; b) Y. Feng & K. Burgess (1999) *Chem. Eur. J.*, **5**, 3261.



12. D.N. Posnett, et al. (1988) *J. Biol. Chem.*, **263**, 1719.
13. S. R. Chhabra, et al. (1998) *Tetrahedron Lett.*, **39**, 1603.
1. <http://scholar.google.co.uk/scholar?cites=7545760316968945982;http://scholar.google.co.uk/scholar?cites=16073846467466700087>
15. N. Ahlborg (1995) *J. Immun. Meth.*, **179**, 269.
16. B. W. Bycroft, et al. in "Peptides, Chemistry, Structure & Biology, Proc. 13th American Peptide Symposium", R.S. Hodges & J.A. Smith (Eds.), ESCOM, Leiden, 1994, pp. 727.
17. D. Lelièvre, et al. (1995) *Tetrahedron Lett.*, **36**, 9317.
18. J. Mack, et al (2001) *J. Pept. Sci.*, **7**, 338.
19. S. K. Sharma, et al. (1999) *J. Pept. Res.*, **53**, 501.
20. P. J. Connolly, et al. (2000) *Tetrahedron Lett.*, **41**, 5187.
21. A. Gopalsamy, et al. (2000) *Bioorg. Med. Chem. Lett.*, **10**, 1715.
22. E. Garanger, et al. (2008) *Chem. Commun.*, 4792.
23. K. S. A. Kumar, et al. (2011) *Bioconjugate Chem.*, **22**, 137.
2. K. Augustyns, et al. (1998) *J. Pept. Res.*, **51**, 127.
25. R. Wilhelm, et al. in "Peptides for the New Millennium, Proc. 16th American Peptide Symposium", G. B. Fields, J. P. Tam & G. Barany (Eds.), Kluwer Academic Publishers, Dordrecht, pp. 58.
26. J. J. Diaz-Mochon, et al. (2004) *Org. Lett.*, **6**, 1127.
27. A. Aletras, et al. (1995) *Int. J. Peptide Protein Res.*, **45**, 488.
28. Z. Flegelova, et al. in "Proc. 32nd European Peptide Symposium", Athens, 2012, poster 43.
29. K. Barlos, Personal communication.
30. L. Bourel, et al. (2000) *J. Pept. Sci.*, **6**, 26
31. Y. Feng & K. Burgess (1999) *Chem. Eur. J.*, **5**, 3261.
32. S. Matysiak, et al. (1998) *Tetrahedron Lett.*, **39**, 1733.
33. F. Albericio, et al (1993) *Tetrahedron Lett.*, **34**, 1549.
3. S. A. Kates, et al. in "Peptides, Chemistry, Structure & Biology, Proc. 13th American Peptide Symposium", R. S. Hodges & J. A. Smith (Eds.), ESCOM, Leiden, 1994, pp. 113.
35. B. Rohwedder, et al. (1998) *Tetrahedron Lett.*, **39**, 1175.
36. P. E. Schneggenburger, et al. (2010) *J. Pept. Sci.*, **16**, 10.
37. M. Meldal, et al. (1997) *Tetrahedron Lett.*, **38**, 2531.
38. J. T. Lundquist & J. C. Pelletier (2001) *Org. Lett.*, **3**, 781.
39. K. Barlos, et al. (1991) *Tetrahedron Lett.*, **32**, 471.
40. K. Barlos, et al. in "Peptides 1992, Proc. 22nd European Peptide Symposium", C. H. Schneider & A. N. Eberle (Eds.), ESCOM, Leiden, 1993, pp. 283.
41. M. P. Coba, et al. (2003) *J. Peptide Res.*, **61**, 17.
42. K. Barlos, et al. (1998) *J. Pept. Res.*, **51**, 19
43. U. Weber & P. Hartter (1970) *Hoppe-Seyler's Z. Physiol. Chem.*, **351**, 138
4. T. M. Postma, et al. (2012) *Org. Lett.*, **14**, 5468.
45. F. Albericio et al. in "Peptides 1992, Proc. 22nd European Peptide Symposium", C. H. Schneider & A. N. Eberle (Eds.), ESCOM, Leiden, 1993, pp. 191.
46. W. Bannwarth, et al. (1992) *Tetrahedron Lett.*, **33**, 4557.
47. N. A. Solé, et al. in "Peptides, Chemistry, Structure & Biology, Proc. 13th American Peptide Symposium", R. S. Hodges & J. A. Smith (Eds.), ESCOM, Leiden, 1994, pp. 93.
48. C. Flouzat, et al. (1997) *Tetrahedron Lett.*, **38**, 1191.
49. W. C. Chan, et al. (1995) *J. Chem. Soc., Chem. Commun.*, 2209.
50. D. J. Evans, et al. Poster 122 presented at the 24th European Peptide Symposium, Edinburgh, 1996.
51. M. Valldosera, et al. (2008) *Int. J. Pept. Res. Ther.*, **14**, 273.
52. K. F. Medzihradsky, et al. (2002) *Lett. Pept. Sci.*, **8**, 1.
53. T. Johnson, et al. (2000) *J. Chem. Soc., Perkin Trans. 1*, 2811.
5. T. Conroy, et al. (2009) *Org. Biomol. Chem.*, **7**, 2255.
55. M. Cudic, et al. in Peptides 2000, Proc. 26th European Peptide Symposium", J. Martinez & J.-A. Fehrentz (Eds), Éditions EDK, Paris, 2001, pp. 203.
56. M. Cudic, et al. (2000) *Tetrahedron Lett.*, **41**, 4527.
57. J. P. Malkinson, et al. (2003) *Org. Lett.*, **5**, 5051.
58. F. Dick, et al. in "Peptides 1996, Proc. of 24th European Peptide Symposium", R. Ramage & R. Epton (Eds.), Mayflower Scientific Ltd., Birmingham, 1998, pp. 339.
59. a) A. S. Dutta (1989) *Chem. Br.*, **25**, 159. ;  
b) K. D. Kopple (1972) *J. Pharm. Sci.*, **61**, 1345.
60. H. F. Brugge, et al. (1994) *Int. J. Peptide Protein Res.*, **43**, 166.
61. a) N. Izumiya, et al. (1981) *Biopolymers*, **20**, 1785. ;b) S. F. Brady, et al. in "Peptides, Structure & Function", V. J. Hruby & D. H. Rich (Eds.), Pierce Chemical Company, Rockford, 1983, pp. 127.
62. a) P. Rovero, et al. (1991) *Tetrahedron Lett.*, **32**, 2639. ;b) J. S. McMurray, et al. (1994) *Pept. Res.*, **7**, 195.
63. J. S. Davies (2003) *J. Pept. Sci.*, **9**, 471.
6. J. Offer, et al. (1996) *J. Chem. Soc., Perkin Trans. 1*, 175.
65. D. Delforge, et al. (1996) *Lett. Pept. Sci.*, **3**, 89.
66. M. Quibell, et al. (1994) *J. Chem. Soc., Chem. Commun.*, 2343.

#### Application 4: Synthesis of cyclo(Tyr-D-Trp-Val-D-Trp-D-Trp-Arg-Glu)

H-Tyr(tBu)-D-Trp-Val-D-Trp-D-Trp-Arg(Pmc)-Glu(O-2-ClTrt resin)-ODMab was prepared automatically using a Perceptive Millipore 9050 peptide synthesizer on 2-ClTrt resin (Figure 4-11). All acylation reactions were carried out using Fmoc-amino acids activated with HBTU (1 eq.) in the presence of DIPEA (2 eq.) and HOBt (1 eq.). Removal of Dmab was effected by flowing 2% hydrazine monohydrate in DMF through the resin bed, until no further release of indazole by-product could be detected by spectrophotometric monitoring at 290 nm. To assess the purity of the intermediate peptide prior to cyclization, a small sample of peptidyl resin was treated with 1% TFA in DCM and the partially protected peptide so released was analyzed by HPLC (Figure 4-12). The resin bound hydrazine salt of glutamic acid was then converted to a DIPEA salt by washing the remainder of the resin with 5% DIPEA in DMF. On-resin cyclization was achieved by treatment of the resin with DIPCDI (1.1 eq.) and HOAt (1.1 eq.) for 18 h. Cleavage was carried out by treatment of the peptidyl resin with TFA/TIS/water/EDT (90:1:5:4) for 2h. The crude peptide was analyzed by HPLC (Figure 13) and PD-MS (Figure 14) [expected M+H<sup>+</sup> 1107.2, found 1107.7].




Fig. 13: HPLC elution profile of crude H-Tyr(tBu)-D-Trp-Val-D-Trp-D-Trp-Arg(Pmc)-Glu-OH.




Fig. 14: HPLC elution profile of crude cyclo(Tyr-D-Trp-Val-D-Trp-D-Trp-Arg-Glu).




Fig. 15: PD-MS of total crude peptide.

## For more information please contact our local offices:

Orders  
Phone 866 645 5476  
Fax 866 645 5439  
Technical Support  
Phone 800 228 9622  
E-mail [technical@novabiochem.com](mailto:technical@novabiochem.com)  
Visit our website  
[www.emdmillipore.com/novabiochem](http://www.emdmillipore.com/novabiochem)

You can also order Novabiochem products through  
our distribution partner, VWR International  
Phone 800 932 5000  
Web [www.vwr.com](http://www.vwr.com)

EMD Millipore  
290 Concord Road  
Billerica MA 01821

CHANGE SERVICE REQUESTED



Product prices and availability are subject to change. Products are warranted only to meet the specifications set forth on their label/packaging and/or certificate of analysis at the time of shipment or for the expressly stated duration. NO OTHER WARRANTY WHETHER EXPRESS, IMPLIED OR BY OPERATION OF LAW IS GRANTED. The products are intended for research purposes only and are not to be used for drug or diagnostic purposes, or for human use. Merck KGaA's products may not be resold or used to manufacture commercial products without the prior written approval of Merck KGaA. All sales are subject to Merck KGaA's complete Terms and Conditions of Sale (or if sold through an affiliated company of Merck KGaA, such affiliated company's complete Terms and Conditions of Sale). NovaSyn® and Novabiochem® are registered trademarks of Merck KGaA in Australia, Germany, Japan, Switzerland, the United Kingdom, and the United States.

©Copyright 2014 Merck KGaA, Darmstadt, Germany. All rights reserved.