

Product Information

Anti-HMGB1 (HMG1), (C-terminal)

produced in rabbit, affinity isolated antibody

Catalog Number **H9539**

Product Description

Anti-HMGB1 (HMG1), (C-Terminal) is produced in rabbit using as immunogen a synthetic peptide corresponding to amino acid 165-180 of human HMGB1 (GeneID: 3146), conjugated to KLH. The corresponding sequence is conserved in mouse and rat. The antibody is affinity-purified using the immunizing peptide immobilized on agarose.

Anti-HMGB1 (HMG1) (C-Terminal) recognizes mouse and rat HMGB1. The antibody may be used in several applications including the detection of HMGB1 by immunoblotting (25 kDa), immunoprecipitation and immunofluorescence. Detection of the HMGB1 band by immunoblotting is specifically inhibited with the immunizing peptide.

The High Mobility Group (HMG) proteins, originally isolated from mammalian cells, were named according to their electrophoretic mobility in polyacrylamide gels. They bind DNA in a non-sequence-specific manner to promote chromatin function and gene regulation. The HMG proteins are subdivided into 3 families based on their functional sequence motif characteristic: the HMGB (formerly HMG-1/-2) family, the HMGN (formerly HMG-14/-17) and the HMGA (formerly HMG-I/Y/C) family. The functional motif of the HMGB family is called the "HMG-box", that of the HMGN family is called the "nucleosomal binding domain;" and that of the HMGA family is called the "AT-hook".¹

The HMGB protein family, including HMGB1, HMGB2, and HMGB3, are highly conserved (more than 80% amino acid identity) and indistinguishable in their biochemical properties. HMGB1 and 2 participate in the regulation of chromatin structure as well as being involved in transcription regulation, DNA repair, recombination, differentiation, and extracellular signaling.²⁻⁷ HMGB1 is a ~ 25 kDa protein of 215 amino acids and consists of two homologous HMG-boxes rich in basic amino acids, and an acidic tail at the carboxy-terminus.¹⁻⁷ Mice that lack HMGB1 die shortly after birth due to hypoglycemia.

They show a defect in transcriptional enhancement of the glucocorticoid receptor that indicates the important role for HMGB1 in proper transcriptional control by specific transcription factors.⁸

Reagent

Supplied as a solution in 0.01 M PBS, pH 7.4, containing 15 mM sodium azide as a preservative.

Antibody concentration: ~1.0 mg/mL

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

Storage/Stability

For continuous use, store at 2-8 °C for up to one month. For extended storage, freeze in working aliquots. Repeated freezing and thawing, or storage in "frost-free" freezers, is not recommended. If slight turbidity occurs upon prolonged storage, clarify the solution by centrifugation before use. Working dilutions should be discarded if not used within 12 hours.

Product Profile

Immunoblotting: a working concentration of 0.5-1 µg/mL is recommended using 3T3 or PC12 cell lysates.

Immunoprecipitation: a working amount of 5-10 µg is recommended using HEK-293T cell lysates.

Note: In order to obtain the best results using various techniques and preparations, we recommend determining the optimal working dilutions by titration.

References

1. Bustin, M., *Trends Biochem. Sci.*, **26**, 152-153 (2001).
2. Bustin, M., and Reeves, R., *Prog. Nucl. Acids Res. Mol. Biol.*, **54**, 35-100 (1996).

3. Onate, S.A., et al., *Mol. Cell. Biol.*, **14**, 3376-3391 (1994).
4. Jayaraman, L., et al., *Genes Dev.*, **12**, 462-472 (1998).
5. Bustin, M., *Mol. Cell. Biol.*, **19**, 5237-5246 (1999).
6. Bianch, M.E., and Beltrame, M., *EMBO Rep.*, **1**, 109-114 (2000).
7. Muller, S., et al., *EMBO J.*, **20**, 4337-4340 (2001).
8. Calogero S., et al., *Nature Genet.*, **22**, 276-280 (1999).

MG,KAA,PHC 08/08-1

Sigma brand products are sold through Sigma-Aldrich, Inc.
Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications.
Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply.
Please see reverse side of the invoice or packing slip.