

69964 MRS Agar (Lactobacillus Agar acc. to De Man, Rogosa and Sharpe)

For the enrichment, cultivation and isolation of all species of *Lactobacillus* from all types of material according to DeMan, Rogosa and Sharpe.

Composition:

Ingredients	Grams/Litre
Universal peptone	10.0
Meat extract	5.0
Yeast extract	5.0
D(+)-Glucose	20.0
Dipotassium hydrogen phosphate	2.0
Triammonium citrate	2.0
Sodium acetate	5.0
Magnesium sulfate (equivalent to 0.2 g/l heptahydrate)	0.1
Manganous sulfate monhydrate	0.05
Agar	12.0
Final pH 6.5 +/- 0.2 at 25°C	

Store prepared media below 8°C, protected from direct light. Store dehydrated powder, in a dry place, in tightly-sealed containers at 2-25°C.

Directions:

Dissolve 61.15 g in distilled water, add 1 ml Tween 80 (Cat. No. P8074) and bring volume to 1000 ml. Boil to dissolve the medium completely. Autoclave at 121°C for 15 minutes.

Incubate the culture up to 3 days at 35°C or up to 5 days at 30°C. If possible, incubate the plates in a CO₂-enriched atmosphere in an anaerobic jar. Do not allow the surface of the plates to dry as this will cause acetate concentration increasing at the surface, which inhibits the growth of lactobacilli.

Principle and Interpretation:

The MRS media formulation was developed by de Man, Rogosa and Sharpe to replace the tomato juice medium and the meat extract tomato juice medium. It is a medium supporting good growth of lactobacilli in general, even those strains which have shown poor growth in existing media, like strains of *L. brevis* and *L. fermenti*. The MRS culture media contain polysorbate (Tween 80), acetate, magnesium and manganese which are known to act as special growth factors for lactobacilli as well as a rich nutrient base. As these media show a very low degree of selectivity, *Pediococcus* and *Leuconostoc* species as well as other secondary bacteria may grow on them. Most of the accompanying microflora can be inhibited by thallium acetate (Cat. No. T8266), sorbic acid (Cat. No. S1626), acetic acid (Cat. No. 33209), sodium nitrite (Cat. No. S2252), cycloheximide (Cat. No. 01810) and polymyxin (Cat. No. P4932). These substances can be used at varying concentrations and combinations, but inevitably a compromise has to be reached between selectivity and productivity of the organism sought.

Cultural characteristics:

Organisms (ATCC)	Growth
<i>Lactobacillus acidophilus</i> (4356)	++
<i>Lactobacillus fermentum</i> (9338)	++
<i>Bifidobacterium bifidum</i> (11863)	++ (anaerobic)
<i>Escherichia coli</i> (25922)	-/+
<i>Pseudomonas aeruginosa</i> (27853)	none to poor



References:

1. J. C. de Man, M. Rogosa and M. Elisabeth Sharpe, Appl. Bact. 23. 130-135 (1960)
2. M. Briggs, J. Dairy Res. 20. 36-40 (1953)
3. G. Reuter, Intern. J. Food Microbiol. 2. 55-68 (1985)
4. ISO/TC 34/SC 6/WG 15, No.3 and 5, Enumeration of Lactobacteriaceae in meat and meat products (1984)
5. Lankaputhra W.E.V., Shah N.P. and Britz M.L. Food Australia 48. 113-118 (1996)
6. W. Hummel, et al., Biocatalysis 2, 293 (1989)
7. L.C. Laleye, et al., Involvement of heterofermentative lactobacilli in development of open texture in cheese, J. Food Prot. 50, 1009 (1987)
8. P. Laloi, et al., Cell-wall-associated proteinase of *L. delbrueckii* subsp. *bulgaricus* CNRZ 397, Appl. Microbiol. Biotechnol. 36, 196 (1991)

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

©2013 Sigma-Aldrich Co. LLC. All rights reserved. SIGMA-ALDRICH is a trademark of Sigma-Aldrich Co. LLC, registered in the US and other countries. Sigma brand products are sold through Sigma-Aldrich, Inc. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see product information on the Sigma-Aldrich website at www.sigmaaldrich.com and/or on the reverse side of the invoice or packing slip.

The vibrant M, Millipore, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. Detailed information on trademarks is available via publicly accessible resources.
© 2018 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the US and Canada.

