

Product Information

Chitinase from Trichoderma viride

Lyophilized powder, ≥600 units/g solid

C8241

Product Description

Enzyme Commission (EC) Number: 3.2.1.14

Synonym: $(1\rightarrow 4)$ -2-acetamido-2-deoxy- β -D-glucan glycanohydrolase

Chitinases are enzymes that catalyze the degradation of chitin. They have been detected in many organisms, including bacteria, fungi, plants, invertebrates, and vertebrates. 1,2 Chitinases are broadly classified as endo- and exochitinases. Endochitinase activity is defined as random cleavage at internal points in the chitin chain. Exochitinase activity is defined as progressive action, starting at the non-reducing end of chitin, with the release of chitobiose or *N*-acetyl-glucosamine units. 2,3 *N*-acetyl-β-glucosaminidase and chitobiosidase are considered exochitinases. The combination of endochitinases and exochitinases results in a synergistic increase in the chitinolytic activity. 4

The chitinolytic enzymes from *Trichoderma viride* are a mixture of extracellular chitinolytic enzymes, 1 which exhibit exo- and endochitinase activities. The major activity was found to be N-acetyl- β -glucosaminidase. The profile of T. *viride* chitinolytic enzymes is similar to that of T. *harzianum*. 5 These activities can be detected after separation by polyacrylamide gel electrophoresis using an agarose overlay containing fluorescent substrates. 2,5

Several theses⁷ and dissertations⁸⁻¹⁰ have cited use of product C8241 in their research.

Precautions and Disclaimer

For R&D use only. Not for drug, household, or other uses. Please consult the Safety Data Sheet for information regarding hazards and safe handling practices.

Product

Specific Activity: ≥ 600 units per g solid

Chitinase Unit Definition: One unit will release 1.0 mg of N-acetyl-D-glucosamine from chitin per hour at pH 6.0 at 25 °C.

N-Acetyl-β-glucosaminidase Unit Definition: One unit will release 1.0 μg of 4-nitrophenol from 4-nitrophenyl N-Acetyl-β-D-glucosaminide per minute at pH 6.1 at 40 °C. 6

The activities of chitobiosidase and of N-acetyl- β -glucosaminidase (NAGase) are determined for each lot. The results are reported on the Certificate of Analysis for each specific lot.

Storage/Stability

The product ships on cooler packs ('wet ice'). The product, as supplied, is stable for 2 years at -20 °C. A solution in 50 mM phosphate buffer, pH 6.1, is stable for at least 5 weeks at 2-8 °C.

Preparation Instructions

The product is soluble in phosphate buffer (pH 6.1) at 1 mg/mL, yielding a clear to faint hazy, tan solution.

One publication reports preparation of stock solutions of this product at 20 mg/mL, in 20 mM sodium phosphate buffer (pH 7).¹¹ Another publication reports preparation of stock solutions of this product at 20 mg/mL, in a buffer of 20 mM sodium citrate with 1 M sorbitol (pH 5.8).¹² We have not tested either condition ourselves.



References

- Rogalski, J. et al., Acta Microbiol. Pol., 46(4), 363- 375 (1997).
- Tronsmo, A., and Harman, G.E., *Anal. Biochem.*, 208(1), 74-79 (1993).
- Felse, P.A., and Panda, T., Bioprocess Engineering, 23(2), 127-134 (2000).
- Bolar, J.P. et al., Transgenic Res., 10(6), 533-543 (2001).
- Haran, S. et al., Mycol. Res., 99(4), 441-446 (1995).
- Frandberg, E., and Schnurer, J., J. Appl. Bacteriol., 76(3), 259-263 (1994).
- Boisseaux, Marion, "Molecular dialogue in the nitrogen-fixing root symbiosis Casuarina/Frankia: characterization of Frankia's symbiotic signal molecules". AgroParisTech (Agrocampus-Ouest, Angers), M.Sc.Eng. thesis, p. 14 (2018).
- Suffrian, Kerstin, "Ion transport and pH homeostasis in coccolithophores".
 Christian-Albrechts-Universität zu Kiel, Dr. rer. nat. dissertation, pp. 38, 51 (2010).
- Ndlovu, Thulile, "Mannoprotein production and wine haze reduction by wine yeast strains". Stellenbosch University, Ph.D. dissertation, p. 122 (2012).
- Brandt, Sophie Charlotte, "Genomische, enzymatische und biotechnologische Charakterisierung einer Pilzsammlung aus Vietnam" ("Genomic, enzymatic and biotechnological characterization of a fungal collection from Vietnam "). Universität Hamburg, Dr. rer. nat. dissertation, p. 77 (2018).
- 11. Scholz, M.J. et al., Eukaryot. Cell, **13(11)**, 1450-1464 (2014).
- 12. Wawra, S. et al., Nat. Comm., 7, 13188 (2016).

Notice

We provide information and advice to our customers on application technologies and regulatory matters to the best of our knowledge and ability, but without obligation or liability. Existing laws and regulations are to be observed in all cases by our customers. This also applies in respect to any rights of third parties. Our information and advice do not relieve our customers of their own responsibility for checking the suitability of our products for the envisaged purpose.

The information in this document is subject to change without notice and should not be construed as a commitment by the manufacturing or selling entity, or an affiliate. We assume no responsibility for any errors that may appear in this document.

Technical Assistance

Visit the tech service page at SigmaAldrich.com/techservice.

Standard Warranty

The applicable warranty for the products listed in this publication may be found at SigmaAldrich.com/terms.

Contact Information

For the location of the office nearest you, go to SigmaAldrich.com/offices.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the U.S. and Canada.

