

Product Information

Papain from papaya latex

Product Number **P 3125**
Storage Temperature 2–8 °C

CAS# 9001-73-4
EC 3.4.22.2
Synonyms: Papainase, papaya peptidase I

Product Description

Papain is a cysteine protease with wide specificity, cleaving peptide bonds of basic amino acids, leucine, or glycine. It also hydrolyzes esters and amides. Papain will digest most protein substrates more extensively than the pancreatic proteases.

Papain consists of a single polypeptide chain with three disulfide bridges and a sulfhydryl group necessary for activity of the enzyme. The specificity of cleavage of the X-Y bond is: where X is a nonspecific amino acid, but arginine and lysine are preferred; and phenylalanine-X-Y bond where residues following phenylalanine are preferred; Y is a nonspecific amino acid residue.¹

Papain is commonly used in cell isolation procedures where it has proven more efficient and less destructive than other proteases on certain tissues. For example, papain has been used to isolate viable, morphologically intact cortical neurons from postnatal rats.² Others have used papain (specifically Product No. P 4762) for the isolation of smooth muscle cells.^{3,4} Papain was found to significantly increase the yield of viable smooth muscle cells while not affecting cell sensitivity to stimulants.⁵

Limited papain digestion has proven useful for structural studies of enzymes and other proteins.⁶⁻⁸ Papain is used in red cell serology to modify the red cell surface to enhance or destroy the reactivity of many red cell antigens as an adjunct to grouping, antibody screening, or antibody identification procedures. Papain has also been shown to be useful in platelet serology.⁹

Papain has also been used in the enzymatic synthesis of amino acids, peptides, and other molecules.¹⁰⁻¹³

Papain is used routinely for the preparation of Fab fragments from IgG. The protease cleaves the antibodies into two Fab fragments, which recognize the antigen specifically with their variable region, and one Fc fragment.¹⁴ IgM may also be digested with papain resulting in high yields of homogeneous Fab preparations.¹⁵

Molecular mass: 23,406 Da (amino acid sequence)¹⁶

Optimal pH: 6.0-7.0

Isoelectric point (pI): 8.75¹⁷; 9.55¹⁸

Spectral properties:

λ_{\max} : 278 nm¹⁹
Extinction coefficient, $E^{1\%} = 25$ ¹⁹
Extinction coefficient, $E^{mM} = 57.6$ (at 280 nm)²⁰

Inhibitors:

Antipain	(Product No. A 6191)
Cystamine	(Product No. C 7255)
Chymostatin	(Product No. C 7268)
Cystatin	(Product No. C 8917)
3,4-Dichloroisocoumarin	(Product No. D 7910)
E-64	(Product No. E 3132)
Ebselen	(Product No. E 3520)
Gly-Gly-Tyr-Arg	(Product No. G 5386)
Leupeptin	(Product No. L 2023)
α_2 -Macroglobulin	(Product No. M 6159)

Potassium halide salts at low pH have also been reported to inhibit papain.¹⁸

Substrates:

pGlu-Phe-Leu p-nitroanilide	(Product No. P 3169)
Gelatin	(Product No. G 2500 or G 8150)

The product is supplied as a buffered aqueous suspension of 2x crystallized papain purified from papaya latex in 0.05 M sodium acetate, pH 4.5, containing 0.01% thymol.

Activity: 16-40 units/mg protein

Unit Definition: One unit will hydrolyze 1.0 μ mole of N- α -benzoyl-L-arginine ethyl ester (BAEE) per minute at pH 6.2 at 25 °C.

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

Preparation Instructions

Immediately prior to use, the enzyme is typically diluted in buffer containing ~5 mM L-cysteine.

Storage/Stability

The product as supplied should be stored at 2–8 °C.

Although papain solutions have good temperature stability, the solution stability is pH dependent. Papain solutions are unstable under acidic conditions, i.e., at pH values below 2.8, there is a significant decrease in activity. For the active enzyme in solution, the loss in activity is about 1-2% per day, probably as a result of autolysis and/or oxidation.

Papain solutions are stable to several denaturing agents, i.e., full activity is maintained after recrystallization in 70% methanol and in 8 M urea solutions. However, there is a significant loss in activity when papain is exposed to 10% trichloroacetic acid or to 6 M guanidine hydrochloride.

A suspension of papain crystals in sodium chloride solution can be kept at 2–8 °C for months without detectable loss in activity. Stabilizing agents include EDTA, cysteine, and dimercaptopropanol.²¹

References

1. Carrey, E.A., in Protein Structure: A Practical Approach, 2nd ed., Creighton, T. E., ed., IRL Press, (New York, NY: 1997), pp 117-144.
2. Huettner, J.E., and Baughman, R.W., J Neurosci., **6**, 3044 (1986).
3. Kinoshita, K., *et al.*, Am. J. Physiol. Gastrointest. Liver Physiol., **285**, G483-G493 (2003).
4. Driska, S.P., *et al.*, J. Appl. Physiol., **86**, 427-435 (1999).
5. Hasegawa, M., *et al.*, Nippon Heikatsukin Gakkai Zasshi, **23**, 35-46 (1987).
6. Margossian, S.S., and Lowey, S., J. Mol. Biol., **74**, 301-311 (1973).
7. Margossian, S.S., and Lowey, S., J. Mol. Biol., **74**, 313-330 (1973).
8. Shiozaki, K., and Yanagida, M., Mol. Cell. Biol., **11**, 6093-6102 (1991).
9. Lown, J.A., and Dale, B.J., Immunohematol., **11**, 140-142 (1995).
10. Xiang, H., *et al.*, Amino Acids, **27**, 101-105 (2004).
11. Rajesh, M., *et al.*, J. Agric. Food Chem., **51**, 2461-2467 (2003).
12. Fukuoka, T., *et al.*, Biomacromolecules, **3**, 768-774 (2002).
13. Burton, S.G., *et al.*, Nature Biotech., **20**, 37-45 (2002).
14. Antibodies: A Laboratory Manual, Harlow, E. and Lane, D., eds., Cold Spring Harbor Laboratory (Cold Spring Harbor, NY, 1988) pp. 628-629.
15. Newkirk, M.M., *et al.*, Hybridoma, **6**, 453-460 (1987).
16. Mitchel, R. E., *et al.*, J. Biol. Chem., **245**, 3485-3492 (1970).
17. J. Biol. Chem., **207**, 533 (1954).
18. Sluyterman, L. A., and de Graaf, M.J., Biochim. Biophys. Acta, **258**, 554-561 (1972).
19. J. Biol. Chem., **236**, 2948 (1961).
20. Pace, C. N., *et al.*, Protein Sci., **4**, 2411-2423 (1995).
21. Arnon, R., Meth. Enzymol., **19**, 226-244 (1970).

ALC,MAM 01/05-1

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.