

Product Information

ANTI-6CKINE, Mouse Developed in Goat, Affinity Isolated Antibody

Product Number **C 6229**

Product Description

Anti-6Ckine is developed in goat using purified recombinant mouse 6Ckine expressed in *E. coli* as immunogen. Affinity isolated antigen specific antibody is obtained from goat anti-6Ckine antiserum by immunospecific purification which removes essentially all goat serum proteins, including immunoglobulins, which do not specifically bind to the peptide.

Anti-6Ckine recognizes recombinant mouse 6Ckine by various immunochemical techniques including immunoblotting, neutralization, and ELISA. The antibody exhibits less than 15 % cross-reactivity with recombinant human 6Ckine.

Mouse 6Ckine is a β - or C-C chemokine identified in the Expressed Sequence Tag (EST) database by three independent groups.^{1,2,3} Known also as Exodus-2 and secondary lymphoid-tissue chemokine (SLC), 6Ckine contains four conserved cysteine (C) residues which are characteristic of β -chemokines.^{1,2,3} Two additional conserved cysteine residues have been found in its unusually long carboxy-terminal domain and consequently the name 6Ckine.^{1,2,3}

Human and mouse 6Ckine are highly conserved and show 86% amino acid homology.¹ Mouse 6Ckine cDNA encodes a 133 amino acid precursor protein, a 23 amino acid signal peptide and a 110 amino acid mature protein.¹ Comparatively, human 6Ckine cDNA encodes a 134 amino acid precursor protein, a 23 amino acid signal peptide and a 111 amino acid mature protein.^{1,3} Recombinant mouse 6Ckine has a predicted molecular mass of approximately 12 kDa (mature protein). In SDS-PAGE under reducing and non-reducing conditions, the recombinant protein migrates with an apparent molecular mass of 16 kDa to 17 kDa.

The expression of mouse 6Ckine has been detected in high levels in the spleen and lung.³ Recombinant mouse 6Ckine is chemotactic *in vitro* for thymocytes and activated T cells.¹ Unlike other C-C chemokines, 6Ckine is not chemotactic for monocytes and

neutrophils.^{1,3} A growing body of work suggests that 6Ckine influences lymphocyte homing to secondary lymphoid organs⁶, integrin-mediated lymphocyte adhesion⁷ and may act via the EBI1 ligand chemokine (ELC) receptor, CCR7^{4,5}.

Reagent

Anti-6Ckine supplied as approximately 100 μ g of antiserum lyophilized from a 0.2 μ m filtered solution in phosphate buffered saline (PBS).

Preparation Instructions

To one vial of lyophilized powder, add 1 ml of sterile phosphate-buffered saline (PBS) to produce a 0.1 mg/ml stock solution of antibody.

Storage/Stability

Prior to reconstitution, store at -20°C . Reconstituted product may be stored at 2° to 8°C for at least one month. For prolonged storage, freeze in working aliquots at -20°C . Avoid repeated freezing and thawing.

Procedure

Anti-6Ckine neutralizes the bioactivity of recombinant mouse 6Ckine. To measure this biological activity, recombinant mouse 6Ckine is incubated with various concentrations of the antibody for 30 minutes at room temperature in a 96 well microplate. Following this preincubation period, 35 μ l of the cytokine-antibody solution (containing recombinant mouse 6Ckine at a final concentration of 0.5 μ g/ml and antibody at concentrations from 0.01 to 10 μ g/ml) is transferred to the lower compartment of a 96 well chemotaxis chamber. The chemotaxis chamber is then assembled using a PVP-free polycarbonate filter (5 micron pore size) and 2×10^6 cells/ well (cultured human lymphocytes) are added to the top chamber. After incubation for 3 hours at 37°C in a 5 % CO_2 humidified incubator, the chamber is disassembled and the cells that have migrated through to the lower chamber are transferred to a working plate and stained using MTT. Absorbance at 540 nm is read on a microplate reader.

The Neutralization Dose₅₀ (ND₅₀) for this antibody is defined as that concentration required to yield one-half maximal inhibition of the 6Ckine activity on a responsive cell line, when 6Ckine is present at a concentration just high enough to elicit a maximum response.

The exact concentration of antibody required to neutralize recombinant mouse 6Ckine activity is dependent on the cytokine concentration, cell type, growth conditions, and the type of activity.

Product Profile

The Neutralization Dose₅₀ (ND₅₀) for this antibody is approximately 8 to 12 µg/ml in the presence of 0.5 µg/ml of recombinant mouse 6Ckine, using cultured lymphocytes in a chemotaxis assay.

For immunoblotting, a working concentration of 0.1 to 0.2 µg/ml detects mouse 6Ckine at approximately 5 ng/lane under non-reducing and reducing conditions.

For ELISAs, a working concentration of 0.5 to 1.0 µg/ml detects recombinant mouse 6Ckine.

Note: In order to obtain best results in different techniques and preparations we recommend determining optimal working dilutions by titration test.

Endotoxin: <10 ng/mg antibody determined by the LAL method.

References

1. Hedrick, J.A. and Zlotnik, A., J. Immunol., **159**, 1589-1593 (1997).
2. Hromas, R. et al., J. Immunol., **159**, 2554-2558 (1997).
3. Nagira, M. et al., J. Biol. Chem., **272**, 19518-19524 (1997).
4. Willmann, K. et al., Eur. J. Immunol., **28**, 2025-2034 (1998).
5. Yoshida, R. et al., J. Biol. Chem., **273**, 7118-7122 (1998).
6. Gunn, M.D. et al., Proc. Natl. Acad. Sci. USA, **95**, 258-263 (1998).
7. Pachynski, R.K. et al., J. Immunol., **161**, 952-956 (1998).

KAA 12/00

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. A dditional terms and conditions may apply. Please see reverse side of the invoice or packing slip.