

63014 Mac Conkey MUG Agar

For the isolation of *Salmonella*, *Shigella* and coliform bacteria, in particular *E. coli*, from diverse material. Bile salts and crystal violet inhibit extensively the gram-positive flora. The presence of lactose and neutral red indicates lactose-positive colonies from which *E. coli* can be identified by UV fluorescence.

Composition:

Ingredients	Grams/Litre
Casein peptone	17.0
Meat peptone	3.0
Sodium chloride	5.0
Lactose	10.0
Bile-salt mixture	1.5
Neutral red	0.03
Crystal violet	0.001
4-Methylumbelliferyl- β -D-glucuronide	0.1
Agar	13.5
Final pH 7.1 +/- 0.2 at 25°C	

Store prepared media below 8°C, protected from direct light. Store dehydrated powder, in a dry place, in tightly-sealed containers at 2-25°C.

Directions:

Dissolve 50.1 g in 1 litre distilled water. Autoclave at 121°C for 15 minutes. Mix well and pour plates. Inoculate the plates as usual after they are solidified. Check the plates under UV light at about 360-370 nm. Slightly blue fluorescence indicates the presence of *E. coli*. For confirmation the indole test can be made with Kovac's reagent (Cat. No. 60983). Cover a colony with 10-20 μ l Kovac's reagent. A change of color to red after 2-10 seconds indicates indole formation.

Principle and Interpretation:

Mac Conkey MUG Agar is used for isolation of *Salmonella*, *Shigella* and coliform bacteria, in particular *Escherichia coli*, from various materials. Casein peptone and Meat peptone provide carbonaceous, nitrogenous and other essential growth nutrients. Most gram positive organisms are inhibited by crystal violet and bile salts. Sodium chloride maintains the osmotic equilibrium. Lactose together with the pH indicator Neutral red are used to detect lactose-positive colonies and *E. coli* can be seen among these due to fluorescence under UV light. Lactose-negative colonies are colourless. Lactose-positive colonies are red and often surrounded by a turbid zone due to the precipitation of bile acids. β -D-glucuronidase, which is produced by *E. coli*, cleaves 4-Methylumbelliferyl- β -D-glucuronide to 4-methylumbelliferone and glucuronide. The fluorogen 4-methylumbelliferone can be detected under a long wavelength UV lamp.



Cultural characteristics after 18-24 hours at 35°C.

Organisms (ATCC)	Growth	Color of colony	Color of medium	Fluorescence	Precipitate
<i>Escherichia coli</i> (25922)	+++	red	red	+	+
<i>Enterobacter aerogenes</i> (13048)	+++	red	red	-	-
<i>Salmonella typhimurium</i> (13311)	+++	colorless	yellowish		-
<i>Salmonella dublin</i> (15480)	+++	colorless	yellowish		-
<i>Shigella sonnei</i> (11060)	+++	colorless	yellowish		-
<i>Proteus mirabilis</i> (29906)	+++	colorless	yellowish		-
<i>Staphylococcus aureus</i> (25923)	-				
<i>Enterococcus hirae</i> (8043)	-				
<i>Bacillus cereus</i> (11778)	-				

References:

1. A. MacConkey, J. Hyg. 8, 333 (1905)
2. Recommendations of the Public Health Laboratory Service Water Subcommittee, J. Hyg. Camb. 51, 268 (1953)
3. European Pharmacopeia II, Chapter VIII, 10
4. A. MacConkey, Bile salt media and their advantages in some bacteriological examinations, J. Hyg., 8; 322 (1908)
5. J. MacFaddin, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. I, Williams and Wilkins, Baltimore (1985)
6. Deutsches Arzneibuch DAB 10
7. A. MacConkey, Bile salt broth, Thompson Yates Lab. Rep., VI/1, 151 (1901)

Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.

The vibrant M, Millipore, and Sigma-Aldrich are trademarks of Merck KGaA, Darmstadt, Germany or its affiliates. Detailed information on trademarks is available via publicly accessible resources.
© 2018 Merck KGaA, Darmstadt, Germany and/or its affiliates. All Rights Reserved.

The life science business of Merck KGaA, Darmstadt, Germany operates as MilliporeSigma in the US and Canada.

